

Shift Work Effect on the Number of Sick Leave among Steel Industry Staffs: A Large Cross-Sectional Study

Shokfeh Maleki¹, Mohammad Gholami Fesharaki^{2*}, Mohsen Rowzati³

¹ Biostatistics MSC student, Biostatistics Department, Faculty of Medical Sciences, TarbiatModares University, Tehran, Islamic Republic of Iran

² Assistant Professor of Biostatistics, Biostatistics Department, Faculty of Medical Sciences, TarbiatModares University, Tehran, Islamic Republic of Iran

³ Physician of search disease, Worksite Follow-Up Unit, Occupational Health Center, Mobarakeh Steel Company, Esfahan, Islamic Republic of Iran

* **Corresponding author:** Mohammad Gholami Fesharaki, Department of Biostatistics, Faculty of Medicine, TarbiatModares University, Ale-Ahmad Highway, Tehran, Iran. Email: Mohammad.gholami@modares.ac.ir

Received 2018 February 24; Accepted 2018 November 10.

Abstract

Background: Sick leave is one of the important areas in Health Professional. Since various unknown factors may be result in sick leave, shift work can be considered as one of the effective factors on the number of sick leave in occupational Medicine.

Objectives: This study aimed to consider the effect of shift work on the number of sick leave among steel industry staffs.

Methods: This cross-sectional study is carried out on Esfahan's Mobarakeh Steel Company (EMSC) staffs in 2016 using census sampling method.

Results: A total number of 21168 staffs with mean age of 39.96 years were entered in this study. 14248 staffs (67.3%) were shift workers. Our results revealed that shift working is significantly correlated with cardiovascular diseases (OR=1.891; $p<0.001$), leukemia (OR=2.29; $p=0.039$), respiratory problems (OR=1.29, $p<0.001$), infectious diseases (OR=1.55, $p<0.001$), mental disorders (OR=1.658, $p<0.001$), and poisonings (OR=0.724, $p<0.001$).

Conclusion: Shift workers are more susceptible to different diseases and subsequently they use higher number of sick leave. Therefore, preventive decisions are needed to reduce the negative effects of shift work.

Keywords: Cardiovascular diseases, Leukemia, Mental disorders, Poisoning, Shift work, Sick leave

1. Background

Shift work, which takes place on a schedule outside the traditional hours, is designed to increase the output of factories and equipment's (1,2).

What is shift work? Shift work is an employment practice designed to make use of, or provide service across, all 24 hours of the clock each day of the week. The practice typically sees the day divided into shifts, set periods of time during which different groups of workers perform their duties. The term "shift work" includes both shifts and day work schedules in which employees change or rotate shifts (3-6).

Recent evidences have revealed that the percentage of shift workers in developed countries is more than 15% (7). Although shift works can help business and technology, they can increase the risk of certain disorders and health problems. This social job can affect human life via different mechanisms (8,9).

Shift work is quite-general in different industries such as steel, oil, petrochemicals and cement industries (7). Over the past few decades many studies considered the effect of shift works on health, immunity, activity and quality of life in shift working staffs (8,10-12). The results indicated the negative impact of shift work on staff's health. Several studies showed a significant correlation between shift work and sick leave (13-16). However, there is no study that considered the relationship between shift work

and sick leave among staffs who work in steel industry in Iran.

2. Methods

This cross sectional study was conducted on workers of Esfahan's Mobarakeh Steel Company (EMSC) in year 2016 using census method sampling method. EMSC is the largest steel producer in Iran country.

Inclusion criteria were official employment with at least two years of work experience; exclusion criteria were dismissal or unwillingness to participate in the study. Medical Ethics Committee of Tarbiat Modares University Medical School approved the study (code number: IR.TMU.REC.1395.398).

In this study shift work defined (regular shift: 2 morning shifts, 2 evening shifts, 2 night shifts and 2 days off, weekly shift: 3 morning shifts, 3 evening shifts, and 1 day off every two weeks, Fridays always off, The morning, evening, and night shifts began at 7 AM, 3 PM, and 11 PM) and day workers worked from 7 AM to 3 PM on weekdays and had Thursdays and Fridays off.

Data Analysis

Data were analyzed using IBM's SPSS Software Version 21. Odds Ratio (OR), Chi-square test was used to compare categorical variables and T-test or Mann-

Whitney U test were used to compare continuous variables. In this study also Logistic regression was used to calculate adjusted OR. In this study, a p -value < 0.05 is considered to be statistically significant.

3. Results

A total number of 21168 staffs were entered in this study, in which 14248 staffs (67.3%) were shift workers. The mean age of workers was 39.96 ± 7.47 years. The basic demographic characteristics of all staffs are summarized in Table 1. The mean number of sick leave for workday and shift work staffs was 12.67 and 12.56 days, respectively. There was no significant difference in the mean number of the sick leave between two groups; however, a significant

difference was observed in mean of work experience between them. Comparison of the odds ratio of daily workers with shift workers and number of sick leave for these workers has been reported (see Table 2). Our results also revealed that shift working is significantly correlated with cardiovascular diseases ($OR=1.891$; $p<0.001$), leukemia ($OR=2.29$; $p=0.039$), respiratory problems ($OR=1.29$, $p<0.001$), infectious diseases ($OR=1.55$, $p<0.001$), mental disorders ($OR=1.658$, $p<0.001$), and poisonings ($OR=0.724$, $p<0.001$). Shift workers suffered from the respiratory problems, infectious disease, bone and joints pains, and poisonings more than workday staffs by 36.09%, 31.41%, 18.03%, and 15.94%, respectively. The mean and SD of the number of sick leave in daily and shift workers is presented in Table 3. According to the

Table 1. the mean \pm SD of age, BMI, work experience and the number of sick leave in daily and shift workers

Variables	Shift				p-value
	Day Worker		Shift Worker		
	Mean	Standard Deviation	Mean	Standard Deviation	
NSD	12.67	15.27	12.56	15.52	0.431
Age	36.68	7.51	39.96	7.47	<0.001
experience	6.544	6.262	8.735	6.085	<0.001
BMI	26.04	3.68	26.07	3.47	0.108

NSD: Number of sick leave days; BMI: Body mass index

Table 2. Comparison of the distribution and the possible chance of sick leave between shift and daily workers

Variables	Total	Shift				Crude		Adjusted		C	
		Day Worker		Shift Worker		OR	P-value	OR	P-value	Lower	Upper
		N	%	N	%						
Visual disturbances(Yes No)	0.02%	2	0.02	2	0.01	0.523	0.611	0.420	0.969	.655	3.247
Diseases of the skin(Yes No)	1.28%	88	1.01	236	1.42	1.408	0.007	1.413	0.015	1.229	2.916
Cardiovascular disease(Yes No)	4.21%	235	2.71	830	5.00	1.891	<0.001	1.97	<0.001	1.068	1.259
Diseases of the joints and bones(Yes No)	31.41%	2664	30.67	5283	31.80	1.054	0.068	1.046	0.175	.692	.798
Leukemia (Yes No)	0.17%	8	0.09	35	0.21	2.29	0.039	1.444	0.378	1.110	1.913
Genital Diseases(Yes No)	7.20%	643	7.40	1178	7.09	0.955	0.37	0.899	0.067	1.44	1.948
Respiratory diseases(Yes No)	36.09%	2797	32.20	6333	38.12	1.297	<0.001	1.429	<0.001	1.850	2.634
Digestive diseases(Yes No)	9.15%	748	8.61	1567	9.43	1.105	0.033	1.027	0.615	1.043	1.327
Disorders of mental organs(Yes No)	6.34%	502	5.78	1103	6.64	1.159	0.008	1.132	0.049	.319	.999
Infectious diseases(Yes No)	15.94%	1276	14.69	2757	16.59	1.155	<0.001	1.318	<0.001	.896	1.120
Endocrine diseases(Yes No)	0.65%	40	0.46	125	0.75	1.638	0.007	1.529	0.059	.960	1.176
Psychopathy(Yes No)	4.62%	285	3.28	885	5.33	1.658	<0.001	1.264	0.003	.986	4.589
Cancer(Yes No)	0.22%	15	0.17	41	0.25	1.43	0.262	1.922	0.102	1.087	1.233
Injuries and poisoning(Yes No)	18.03%	1824	21.00	2737	16.47	0.742	<0.001	0.762	<0.001	1.259	1.425

Table 3. the mean \pm SD of the number of sick leave in daily and shift workers based on diseases type

Variables	Shift				p-value
	Day Worker		Shift Worker		
	Mean	Standard Deviation	Mean	Standard Deviation	
Visual disturbances	0.029	.02	0.000	0.00	0.042
Diseases of the skin	1.185	.13	1.818	.18	0.008
Cardiovascular disease	3.121	.24	8.219	.46	<0.001
Diseases of the joints and bones	41.821	.85	51.207	1.02	<0.001
Leukemia	0.130	.04	0.190	.05	0.373
Genital Diseases	9.205	.41	9.833	.45	0.330
Respiratory diseases	39.624	.72	55.131	1.00	<0.001
Digestive diseases	10.173	.37	11.567	.42	0.018
Disorders of mental organs	6.633	.29	8.640	.43	<0.001
Infectious diseases	16.243	.46	19.448	.52	<0.001
Endocrine diseases	0.419	.07	1.053	.16	0.002
Psychopathy	4.104	.25	7.819	.45	<0.001
Cancer	0.159	.05	0.618	.26	0.148
Injuries and poisoning	27.182	.58	22.487	.56	<0.001

information reported in this table we can say except , leukemia , genital and cancer diseases, the mean of sick leave is higher in shift worker rather than day worker.

4. Discussion

A great number of studies showed that shift working is correlated with sleep failure, stress and decreased activity of persons. On the other hand, organizations have recently focused on processes and subjects that affect the cost and efficiency. One of these subjects is staffs sick leave. In this study, we investigated the effect of shift work on the number of sick leave among male staffs who work at Esfahan's Mobarakeh Steel Company. Different parameters such as age, work experiences, BMI, and type of diseases were evaluated. Our data showed that shift work is significantly associated with increased risk of cutaneous, cardiovascular, pulmonary, infectious diseases, mental disorders, disturbance of the sensory organs, and poisonings. Our results were in agreement with a study by Merkus et al (13). Although we could not find a significant difference in the mean number of sick leave between workday and shift work staffs, the distribution of sick leave among shift workers was significantly higher than that in daily workers. We also found that shift workers are more susceptible to cutaneous, cardiovascular, leukemia as well as, digestive problems, glands failure, mental disease, cancer and job injuries. Since shift workers are usually healthier than daily workers, their disease is more dangerous compared to the daily workers. For this reason, non-significant difference was observed in the mean number of sick leave between two groups.

This study also demonstrated that the chance of chronic diseases and as the result sick leave among shift workers is higher than daily workers. Since shift work staffs work higher than daily workers, they are more exposed to the workplace pollutions and tensions (6). Our findings have also revealed that age and work experiences have significant influence on the number of sick leave. Older individuals have the increased risk of different diseases and as the result they have to use greater number of sick leave. This result is in agreement with the findings of other previous studies (6,17,18).

In this research, the risk of cardiovascular disease in shift worker was greater (by 97%) than that in daily workers. Furthermore, the risk of pulmonary and infectious diseases in shift workers was higher by 42% and 31%, respectively. Additionally, the chance of mental disorders and poisonings in shift work staffs was 26% and 0.76% greater than daily workers. These findings were comparable with the results of studies provided by Pallesen et al (16), Norder et al (19), Oenning et al (15), Koller et al (20) and Merkus et al (13), that showed a significant effect

of shift work on the number of sick leave. The strength of this study is related to its novelty and large sample size. In this study, all participants were men who working at steel industry. Since steel company is considered as a heavy industry, the results may be limited for this industry that is a limitation of the study.

5. Conclusion

According to the result of this study, shift workers are more susceptible to different diseases and subsequently they use higher number of sick leave. Therefore, preventive decisions are needed to reduce the negative effects of shift work.

Acknowledgments

The authors gratefully acknowledge financial support from TarbiatModares University and wish to thank all the personnel, especially the staff of Industrial Medicine Department of EMSC, for their cooperation throughout the study.

Conflicts of interest

In this study we have not any conflicts of interest

References

1. Tayyari F, Smith JL. Occupational ergonomics: principles and applications. London: Chapman & Hall; 1997.
2. Fesharaki GM, Rozati M, Tanhai A. The longitudinal study of the relationship between work shift and blood pressure in workers of Mobarakeh Steel Company of Isfahan in 2007-2009. *Arak Med Univ J*. 2010;13(4):68-74.
3. Gholami-Fesharaki M, Kazemnejad A, Zayeri F, Sanati J, Akbari H. A retrospective cohort study on factors associated blood pressure using multilevel modeling. *ARYA Atheroscler*. 2013;9(5):293-9. [PubMed: 24302938].
4. Gholami-Fesharaki M, Kazemnejad A, Zayeri F, Rowzati M, Sanati J, Akbari H. Multicenter historical cohort study of the relationship between shift work and blood pressure. *ARYA Atheroscler*. 2014;10(6):287-91. [PubMed: 25815017].
5. Gholami Fesharaki M, Kazemnejad A, Zayeri F, Sanati J, Akbari H. Historical cohort study on the factors affecting blood pressure in workers of polyacryl iran corporation using bayesian multilevel modeling with skew T distribution. *Iran Red Crescent Med J*. 2013;15(5):418-23. doi: 10.5812/ircmj.10930. [PubMed: 24349731].
6. Staff who work long shifts more likely to call in sick, according to new research. *Nurs Manag*. 2017;24(2):7. doi: 10.7748/nm.24.2.7.s6. [PubMed: 28446094].
7. Fischer FM. What do petrochemical workers, healthcare workers, and truck drivers have in common? Evaluation of sleep and alertness in Brazilian shiftworkers. *Cad Saude Publica*. 2004;20(6):1732-8. doi: /S0102-311X2004000600033. [PubMed: 15608877].
8. Choobineh A, Soltanzadeh A, Tabatabaie SH, Jahangiri M. Shift work-related health problems among petrochemical industries employees. *Sci Med J*. 2011;11:141-51.
9. Repetti RL. Short-term and long-term processes linking job stressors to father-child interaction. *Soc Dev*. 1994;3(1):1-15. doi: 10.1111/j.1467-9507.1994.tb00020.x.
10. Demerouti E, Bakker AB, Bulters AJ. The loss spiral of work pressure, work-home interference and exhaustion: reciprocal

- relations in a three-wave study. *J Vocational Behav.* 2004;**64**(1):131-49. doi: [10.1016/S0001-8791\(03\)00030-7](https://doi.org/10.1016/S0001-8791(03)00030-7).
11. Kaliterna LL, Prizmic LZ, Zganec N. Quality of life, life satisfaction and happiness in shift-and non-shiftworkers. *Rev Saude Publica.* 2004;**38**:3-10. doi: [10.1590/s0034-89102004000700002](https://doi.org/10.1590/s0034-89102004000700002). [PubMed: [15608908](https://pubmed.ncbi.nlm.nih.gov/15608908/)].
 12. Janssen D, Nachreiner F. Health and psychosocial effects of flexible working hours. *Rev Saude Publica.* 2004;**38**:11-8. doi: [10.1590/s0034-89102004000700003](https://doi.org/10.1590/s0034-89102004000700003). [PubMed: [15608909](https://pubmed.ncbi.nlm.nih.gov/15608909/)].
 13. Merkus SL, van Drongelen A, Holte KA, Labriola M, Lund T, van Mechelen W, et al. The association between shift work and sick leave: a systematic review. *Occup Environ Med.* 2012;**69**(10):701-12. doi: [10.1136/oemed-2011-100488](https://doi.org/10.1136/oemed-2011-100488). [PubMed: [22767871](https://pubmed.ncbi.nlm.nih.gov/22767871/)].
 14. van Drongelen A, Boot CR, Hlobil H, van der Beek AJ, Smid T. Cumulative exposure to shift work and sickness absence: associations in a five-year historic cohort. *BMC Public Health.* 2017;**17**(1):67. doi: [10.1186/s12889-016-3906-z](https://doi.org/10.1186/s12889-016-3906-z). [PubMed: [28077111](https://pubmed.ncbi.nlm.nih.gov/28077111/)].
 15. Oenning NS, Carvalho FM, Lima VM. Risk factors for absenteeism due to sick leave in the petroleum industry. *Rev Saude Publica.* 2014;**48**(1):103-22. doi: [10.1590/s0034-8910.2014048004609](https://doi.org/10.1590/s0034-8910.2014048004609). [PubMed: [24789643](https://pubmed.ncbi.nlm.nih.gov/24789643/)].
 16. Vedaa O, Pallesen S, Waage S, Bjorvatn B, Sivertsen B, Erevik E, et al. Short rest between shift intervals increases the risk of sick leave: a prospective registry study. *Occup Environ Med.* 2017;**74**(7):496-501. doi: [10.1136/oemed-2016-103920](https://doi.org/10.1136/oemed-2016-103920). [PubMed: [27827302](https://pubmed.ncbi.nlm.nih.gov/27827302/)].
 17. Hubertsson J, Petersson IF, Thorstensson CA, Englund M. Risk of sick leave and disability pension in working-age women and men with knee osteoarthritis. *Ann Rheum Dis.* 2013;**72**(3):401-5. doi: [10.1136/annrheumdis-2012-201472](https://doi.org/10.1136/annrheumdis-2012-201472). [PubMed: [22679305](https://pubmed.ncbi.nlm.nih.gov/22679305/)].
 18. Pouryaghoub G, Mehrdad R, Rafiee SF, Mahmoodi F. Risk factors of sickness absence. *Occup Med.* 2016;**8**(3):21-30.
 19. Norder G, Roelen CA, Bültmann U, van der Klink JJ. Shift work and mental health sickness absence: a 10-year observational cohort study among male production workers. *Scand J Work Environ Health.* 2015;**41**(4):413-6. doi: [10.5271/sjweh.3501](https://doi.org/10.5271/sjweh.3501). [PubMed: [25945780](https://pubmed.ncbi.nlm.nih.gov/25945780/)].
 20. Koller M. Health risks related to shift work. An example of time-contingent effects of long-term stress. *Int Arch Occup Environ Health.* 1983;**53**(1):59-75. [PubMed: [6654503](https://pubmed.ncbi.nlm.nih.gov/6654503/)].