

Application Validation Methods in EHR Evaluation Models: A Systematic Literature Review

Zahra Ebnehoseini¹, Marziyhe Meraji², Hamed Tabesh^{3,*}

¹Health Information Technology unit of Economic Health Department, Mashhad University of Medical Sciences, Mashhad, Iran

²Department of Health Information Technology, School of Paramedical Sciences, Mashhad University of Medical Sciences, Mashhad, Iran

³Department of Medical Informatics, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

* **Corresponding author:** Hamed Tabesh. Associate Professor of Biostatistics, Department of Medical Informatics, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran. Email: TabeshH@mums.ac.ir.

Received 2023 February 28; Accepted 2023 May 16.

Abstract

Background: Over the last few decades, several theoretical frameworks have been proposed to evaluate electronic health records (EHRs). These frameworks provide a theoretical basis for assessing the impact and outcomes of technology adoption in healthcare settings. This can help identify areas for improvement and ensure that EHRs effectively support healthcare delivery and patient care.

Objectives: The purpose of this study is to present a comprehensive review of the use of validation methods in Electronic health records.

Methods: Out of a total of 62 EHR evaluation frameworks in our previous literature review, at the final stage, 34 relevant articles were included for analysis. Variables such as participants and study setting, analysis software, data gathering methods, missing data, and outlier handling, theoretical basis models to develop the EHR evaluation model, the relationship between variables of the EHR evaluation models with evaluation items, sampling technique and sample size reliability assessment methods and values, and statistical validation methods and criteria values were extracted.

Results: Among the 34 papers that disclosed the validation methods utilized, the most widely used technique was Structural Equation Modeling (SEM), employed in 26.5% of the studies. Other methods utilized were Confirmatory Factor Analysis (CFA), and Exploratory Factor Analysis (EFA). A reliability assessment was performed in 82% of the articles. Cronbach's alpha and composite reliability (CR) were popular reliability (internal validation) methods on identified papers.

Conclusion: It is our belief that the results of this study can assist researchers in examining and modifying EHR evaluation frameworks to suit their specific needs. Additionally, we believe that our findings serve as a solid foundation for the creation of new EHR evaluation frameworks. Furthermore, we recommend that researchers utilize the findings presented in this article to enhance the implementation and utilization of SEM, CFA, and EFA methods in EHR evaluation models.

Keywords: Electronic medical records, Electronic health records, Evaluation frameworks, Evaluation models, Evaluation theory

1. Background

In recent years, the use of electronic health records (EHRs) has dramatically increased among healthcare organizations across the globe. Although the assessment of the quality and performance of the systems is a significant concern (1), there is a need for an evaluation framework for EHRs. This framework can be adopted or developed to ensure that standard evaluation methods are followed (2). Over the last three decades, various theoretical frameworks have been suggested to evaluate and clarify the adoption and behaviors related to introducing Information and Communication Technology (ICT). Some measures have been developed to evaluate how effectively a technology aligns with the tasks of its users, and these instruments have been validated to assess the task-technology fit (3).

Electronic health records are implemented in complex healthcare settings, and their success depends on a wide array of interacting factors. Multiple studies have evaluated the success and failure factors of electronic health records and found that all influential variables do not have equal effects on system success and can interact with each other. Therefore, it is crucial to address all these factors to ensure the successful implementation and adoption of electronic health

records in healthcare centers. The EHR evaluation methods have assessed the factors influencing system success and failure. The accuracy of these models is a significant issue.

2. Objectives

The main aim of this study was to address two crucial questions: "What is the validation method used for evaluation models?" and "What are the various variables present in the cause-and-effect models of EHR evaluation?" The primary objective of this paper is to conduct a comprehensive review of validation methods used in EHR evaluation models. Previous studies have demonstrated that review articles are useful resources for researchers seeking quick and easy access to relevant information.

3. Methods

A systematic search of English literature from January 2007 to August 2017 was conducted in PubMed, Scopus, ScienceDirect, and Cochrane databases to identify evaluation frameworks for electronic health records (4).

Criteria for considering studies

A total of 8,276 records were retrieved, and 62 studies met the inclusion criteria (4). In this study, we used the results of our previous systematic review. In the current review, among 62 articles, studies that propose an EHR evaluation model were included in the analysis. On the other hand, the exclusion criteria were as follows: 1. The use of existing evaluation framework for EHR evaluation without changes, 2. Investigating the relationship between contributed factors based on the existing evaluation framework, and 3. No reporting valuation methods and values or validity assessment.

Eligibility screening and data extraction

Data were extracted by the first author (ZE) and supervised by a second author (HT) using a spreadsheet file. A random sample of seven included papers (one-third of the total included papers) that used Structural equation modeling (SEM), Exploratory Factor Analysis (EFA), Confirmatory Factor Analysis (CFA), and other validation methods were selected for piloting the data extraction using a separate spreadsheet file.

Data synthesis

The extraction checklist was modified as necessary during the study. Data extraction included for the paper included:

- Participants and study setting
- Analysis software
- Data gathering methods, missing data, and outlier handling
- Theoretical basis models to develop the EHR evaluation model
- Relationship between variables of the EHR evaluation models with evaluation items (independent, moderating, dependent items)
 - Sampling technique and sample size
 - Reliability assessment methods and values

To assess the reliability and validity of the measurement scales used in the studies, Cronbach's' alpha and Composite Reliability (CR) were extracted from the papers. The mean and standard variation for these variables were calculated.

- Statistical validation methods and criteria values

Structural equation modeling approach

The articles that used the SEM approach for validation methods following statistical criteria were extracted. A list of Fit indices was prepared based on previous studies as follows:

Chi-square (χ^2), P-value, Degree of freedom (df), χ^2/df , Delta Chi-square, Comparative Fit Index

(CFI), Differences in CFI, Incremental Fit Index (IFI), Non-Normed Fit Index (NNFI), Normed Fit Index (NFI), Parsimony Comparative Fit Index (PCFI), Parsimony Normed Fit Index (PNFI), Tucker-Lewis Index (TLI), Relative Fit Index (RFI), Goodness-of-Fit Index (GFI), Adjusted Goodness-of-Fit Index (AGFI), Parsimony Goodness-of-Fit Index (PGFI), Full Information Maximum Likelihood with Missing Data (FIMIN), Noncentrality parameter (NCP), Root Mean Square Residual (RMR), Standardized Root Mean Square Residual (SRMR), Root Mean Square Error of Approximation (RMSEA), RMSEA with a confidence interval, P-value for Test of Close Fit (PCLOSE), Akaike Information Criterion (AIC), Consistent Akaike Information Criterion (CAIC), Browne-Cudeck Criterion (BCC), Bayesian Information Criterion (BIC), Expected Cross Validation Index (ECVI), and Modified Expected Cross Validation Index (MECVI) (5-7). The mean and standard deviation were not calculated for the criteria that were affected by sample size, and only the frequency and range (minimum and maximum values) were reported in the studies.

Exploratory Factor Analysis and Confirmatory Factor Analysis

KMO (Kaiser-Meyer-Olkin) value and Barlett test p-value were reported in the EFA approach, and Average Variance Extracted (AVE) values were extracted for the CFA approach.

Regression and correlation approach

Since the current study did not specifically investigate the relationship between variables, coefficients, and p-values were not extracted in studies that used correlation and regression for evaluating EHR models. Moreover, articles that did not report Cronbach's alpha and only conducted correlation or regression analysis were excluded from the current study.

4. Result

According to our previous study, a total of 64 full-text papers were included in the current study. Nonetheless, during the first screening, 21 articles were excluded since they did not propose an EHR evaluation model and instead adopted an existing evaluation framework without modification ([Figure 1](#)). Among the remaining 43 articles, three papers developed an EHR evaluation model; nonetheless, the validation method was not performed. In addition, four articles did not report the values of the validation method used. The study findings are summarized in [Tables 1](#) and [Table 2](#).




Figure 1. Flow chart of the study selection process

Table 1. Evaluation items in EHR evaluation models

Row	Authors	Basis for evaluation framework development	Evaluation items			Diagram of EHR Evaluation Model	Data gathering methods	Analysis software	Sampling technique	Missing data	Outliers
			Independent	Moderating	Dependent						
505	Lambooj et al. (8)	Literature review.	Authentic leadership\ support of administrative department\ Support of IT department\ support of HR department\ Bottom up communication in the hospital\ open culture in hospital \ Innovative culture in hospital	Ease of use\ alignment of EMR with daily routine\ perceived added value\ timeliness of use	Perceived quality of patient data	Yes	In-person	Not reported	Convenience sampling	No	No
601	Mohd Salleh et al. (9)	ISSM, EHR System Effectiveness Model.	Adequate infrastructure\ system interoperability\ perceived security control\ system compatibility\	Did not have	Provider performance	Yes	In-person	Smart PLS	Convenience sampling	No	No
504	Kim et al. (End-User Acceptance Model) (10)	TAM, UTAUT.	Performance expectancy \effort expectancy \social influence \facilitating conditions \	Attitude	Behavior intention to use	Yes	Online	SAS-AMOS	not report	No	No
2	Liu et al. (11)	TAM, DFM.	perceived mobility	Perceived usefulness\ perceived ease of use\ perceived threat	Behavior intention	Yes	In-person	Not reported	Convenience sampling	No	No
507	Sintonen et al. (12)	TAM, TBP.	Complexity\ reliability	Perceived usefulness \behavioral control\ usage intention	Actual use	Yes	Email	Smart PLS	Convenience sampling	No	No
512	Gan et al. (13)	TTF, SCT.	Organizational perceived task technology fit	Organizational valence to EHR\ social contagion	Intention to adopt EHR	Yes	In-person	not reported	Convenience sampling	Yes	No
63	Kralj et al. (Initial Framework Model) (14)	Literature review.	Business functionality\ privacy and data security\ domain (health) functionality\ data exchange with patient\ ergonomic functionality \additional services	*	*	Yes	Online	SPSS	not report	No	No
68	Wang et al. (User Acceptance Model of EPR) (15)	TAM	Personal characteristics: type of user, gender, age, education, occupation, number of outpatient visit\ work characteristics: job seniority, level of position, title of physician, level of hospital\ human aspects: reason to use EMR, level concerns\ technology	Did not have	Behavioral intention of use: attitude to privacy attitude to hospital information support	Yes	Online	SAS	Convenience sampling	No	No

40	Michel-Verkerke et al. (16)	IT-Used model	aspects: usefulness, major information source Task support satisfaction\ interface satisfaction\ compatibility\ collaboration\ learnability\ ease of use\ support of use	*	*	No	Online	Not reported	not report	No	No
519	Hysong et al. (17)	UTAUT, Job JDRM.	Monitoring and feedback\ provider\ supportive norm\ provider perceptions of value\ training	Physician satisfaction\ intention to quit	Voluntary turn over	Yes	Online	SPSS-AMOS	Convenience sampling	No	No
48	Gagnon et al. (18)	TAM, TIB, DFM.	Perceived consequences\ facilitating conditions\ computer self-efficacy\ demonstrability of results\ personal identity\ social norm\ professional norm\ resistance to change	Perceived usefulness\ perceived ease of use	Behavioral intention to use	Yes	Online	AMOS	Convenience sampling	Yes	No
545	Iqbal et al. (19)	TAM, UTAUT, TPB.	Computer Self-efficacy\ Subjective Norm\ Security\ Privacy\	Perceived usefulness\ Perceived ease to use	Intention to use	Yes	Online	SPSS	Convenience sampling	No	No
548	Kuo et al. (20)	TAM, TRI.	Optimism\ innovativeness\ insecurity\ discomfort	Perceived Ease Of Use\ Perceived Usefulness	Behavioral intention to use	No	Unclear	Smart PLS	bootstrapping resampling	No	No
14	Lu et al. (21)	TAM, ISSM.	System quality\ information quality\ service quality	Perceived ease of use\ perceived usefulness	Hospital information acceptance	Yes	In-person	LISREL-SPSS-STATA	simple random sampling	No	No
21	Aggelidis et al. (22)	ISSM, Bailey and Pearson's model, Doll and Torkzadeh.	Information quality\ system quality	Insourcing support\ outsourcing support	Overall satisfaction	Yes	In-person	AMOS	Convenience sampling	No	No
20	Chen and Hsiao (23)	TAM, HOT-FIT.	Human characteristic: self-efficacy, compatibility\ organizational characteristic: top management support, project team competency \ technology characteristic: system quality, information quality	Perceived usefulness\ perceived ease of use	Hospital information system acceptance	No	In-person	AMOS	not report	No	No
521	Lin et al. (24)	TAM, Lin et al., Bhattacharjee and Hikmet's.	Perceived threat	Perceived usefulness\ perceived inequity	Behavior intention	Yes	In-person	SPSS-AMOS	Convenience sampling	No	No
122	Carayon et al. (Model of electronic health records (EHR) acceptance) (25)	Chin et al., Carayon et al., Hoonakker et al.	Usefulness\ Usability	*	HIS acceptance	No	In-person	Not reported	Convenience sampling	Yes	No
109	Tilahun et al. (26)	ISSM	System quality\ information quality\ service quality\ computer literacy\	User satisfaction\ system use	Perceived net-benefit	Yes	In-person	SPSS-AMOS	simple random sampling	No	No
506	Steininger et al. (27)	TAM	Privacy concerns\ social influence\ HIT experience\ cost saving\ stakeholder benefit\ improvement	Perceived usefulness	Intention to use	Yes	In-person or online	Not reported	not report	No	No
32	Kowitlawakul et al. (28)	TAM	Self-efficacy	Perceived ease of use\ behavioral intension\ attitude toward using	Intention to use	No	Online	SPSS-AMOS	Convenience sampling	Yes	Yes
511	Hsieh et al. (29)	TBP	Perceived usefulness\ perceived ease of use\ compatibility\ interpersonal influence\ governmental influence\ self-efficacy\ facilitating conditions\ situational normality\ structural assurance	Attitude \ subjective norm\ perceived behavior control\ institutional trust\ perceived risk	Usage intention	No	Unclear	Not reported	not report	No	No
537	Abdekhoda et al. (30)	TAM	Management support \adequate training \physicians' involvement \doctor-patient relationship	Perceived usefulness\ perceived ease of use	System usage	Yes	In-person	AMOS	Convenience sampling	No	No
7	Messeri et al. (31)	ISSM	System Quality\ IT support	User Satisfaction: ease of use\satisfaction	Individual Impact: productivity\ prevention\ decision making, workflow\	Yes	In-person	STAT	Convenience sampling	Yes	No

56	Hsu et al. (32)	DOI	Relative advantage\ Compatibility\ Complexity\ Trialability\ Observability\ Unit\ Seniority	Did not have	Behavior intention	Yes	Online	Not reported	Convenience sampling	No	No
53	Leblanc et al. (33)	TPB	External variables: age, gender, education, prior use of EHR, type of organization	Behavioral beliefs\ Normative beliefs\ facilitating conditions\ Attitude\ subjective norm\ perception of behavioral control	Intention	Yes	In-person	Not reported	Convenience sampling	No	No
50	Chen and Hsiao (34)	TAM	System quality\ Information quality\ Service quality	Perceived usefulness\ Perceived ease of use\	HIS acceptance	Yes	In-person	Not reported	Convenience sampling	No	No
26	Hsiao et al. (35)	TAM	IS construct\ personal construct\ organizational construct\ Perceived ease of use\	Did not have	Hospital information system acceptance	Yes	In-person or online	SPSS	Convenience sampling	No	No
33	Sicotte et al. (36)	ISSM	Qualities of PACS\ data quality\ quality of technical support	Overall satisfaction\ Use	Future intention	Yes	In-person or online	Not reported	Convenience sampling	No	No
602	Erlirianto et al. (37)	HOT-Fit	Technology:(system quality\ information quality\ service quality)	Human: system use, user satisfaction\ Organization: structure, environment	Net benefit	Yes	Unclear	GeSCA	Convenience sampling	No	No
11	Ho et al. (38)	IPA	Installation and Maintenance \ Product Effectiveness \ System Function \ Customer Service \ Personal Information\	*	*	Yes	Unclear	Not reported	Convenience sampling	Yes	No
52	Devine et al. (39)	ITAM	Finesse	Perceived ease of use\ Perceived usefulness	Intent to use	Yes	In-person or online	STAT	Convenience sampling	No	No
64	Otieno et al. (40)	ISSM	EMR use\ Quality of EMR systems\ User satisfaction	*	*	Yes	Online	SPSS	Convenience sampling	No	No
66	Otieno et al. (41)	ISSM	EMR use\ Quality of EMR systems\ User satisfaction	*	*	Yes	Online	SPSS	Convenience sampling	Yes	No

Star sings (*) shows unclear causal relationship between variables
 Ammenwerth and Dekeizer (ADK), Computer Anxiety Model (CAM), Diffusion of innovation (DOI), Composite Index (CI)Dual Factor Model (DFM), Fit Between Individuals, Task and Technology framework (FIT), Human, Organization And Technology-Fit Factors (HOT-Fit), Information Success Model (ISSM), Job Demands Resource Model (JDRM), Theory of Planned Behavior (TPB), Technology Acceptance Model (TAM), Task and Technology and Fit (TTF),Technology Readiness Index (TRI), Unified Theory of Acceptance and Use of Technology (UTAUT).

Participants and study setting

The majority of the studies (n=23; 67.64%) were conducted in hospitals, followed by primary care clinics (n=7; 20.58%). Other locations included

physician offices, specialty associations, and universities. In 64% of the studies (n=22), physicians participated in the evaluation of EHR, while in 56% of the studies (n=19), nurses were involved.

Table 2. The validation models and values with study characteristics of EHR evaluation models

	Authors	Validation methods	SEM, CFA, EFA values	Reliability methods and values	Sample Size	Participants	Study setting
1	Lambooij et al. (8)	505 SEM	χ^2 chi2= 3852.62, CFI = 0.93, TLI = 0.93, RMSEA = 0.046	Cronbach's alpha: 0.70 to 0.92	914	Physicians\nurse	Hospital
2	Mohd Salleh et al. (9)	601 CFA	AVE= 0.583-0.823	CR: 0.807-0.926	367	Nurses\resident\medical officer	Hospital
	Kim et al. (End-User Acceptance Model) (10)	504 SEM	χ^2 =449.217, df=108, P-value=0.000, TLI=0.910, CFI=0.936, and RMSEA=0.84	Cronbach's alpha: 0.844 to 0.979	942	Physicians and nurses	Hospital
3	Liu et al. (11)	2 CFA	AVE= 0.81-0.88	Cronbach's alpha: 0.89 to 0.96\ CR: 0.93-0.97	158	Physicians	Hospital
4	Sintonen et al. (12)	507 CFA	AVE= 0.762-1.000	CR: 0.862 to 1.000	187	Physicians and head nurses	Primary Care Clinic
5	Gan et al. (13)	512 CFA	AVE= 0.511-0.886	CR: 0.733 to 0.955	51	Health organization management students	University
6	Kralj et al. (Initial Framework Model) (14)	63 EFA	KMO= 0.857 -0.962, Bartlett's test: P-value < 0.001, factor loading:****	Cronbach's alpha: 0.637 to 0.973	384	Physicians and other non-clinical participants	A Medical Association
7	Wang et al. (User Acceptance Model of EPR) (15)	68 EFA	KMO=0.46, 0.63, and 0.48 physicians, for MRS, and patients, respectively	Cronbach's alpha: 0.72 to 0.85	379	Physicians, non-clinical participants, and patient	Hospital
8	Michel-Verkerke et al. (16)	40 EFA	KMO=0.910, Bartlett's test: (P-value < .001)	Cronbach's alpha: 0.896, 0.903, and 0.907	222	Physicians, nurses, and other clinicians	Hospital
9	Hysong et al. (17)	519 SEM	RMSEA= 0.04, PCLOSE = 0.47	not reported	2590	Physicians	Medical Facilities

10	Gagnon et al. (18)	48	SEM	CFI=0.94, IFI=0.94, TLI= 0.92	Cronbach's alpha: 0.66 to .91	157	Physicians	A Medical Association
11	Iqbal et al. (19)	545	Correlation analysis	*****	Cronbach's alpha: > 0.90	1097	Primary care clinic, physicians	Hospital
12	Kuo et al. (20)	548	CFA	AVE=0.69-0.87	Cronbach's alpha: =0.77 to 0.95 CR= 0.87 to 0.96	878	Nurses	Hospital
13	Lu et al. (21)	14	CFA-SEM	AVE 0.80-0.90, $\chi^2 = 3294.48$, GFI=0.68, PGFI = 0.62, NFI= 0.98, NNFI= 0.98, CFI=0.98, RMSEA=0.08, RMR=0.035, SRMR=0.048.	Cronbach's alpha: 0.90 to 0.98, CR= 0.74 to 0.87	297	Nurses	Hospital
14	Aggelidis et al. (22)	21	SEM	$\chi^2/Df= 1.82$, CFI=0.015, GFI=0.958, NFI=0.928, RMR=0.928, 0.980, RMSEA= 0.064	Did not reported	341	Physicians, nurses, other clinicians, and other non-clinical participants	Hospital
15	Chen and Hsiao (23)	20	CFA-SEM	AVE=0.756 to 0.933, $\chi^2 = 1124.56$, d.f.=466.00, $\chi^2/d.f.=2.41$, GFI=0.85, NFI=0.90, NNFI=0.91, IFI=0.93, CFI=0.90, RMSEA)=0.08	Cronbach's alpha: 0.701 to 0.908, CR:0.857 to 0.967	202	Physicians	Hospital
16	Lin et al. (24)	521	CFA-SEM	AVE=0.76 to 0.84, $\chi^2 = 1124.56$, d.f.=466.00, $\chi^2/d.f.=1.13$, GFI=0.93, NFI)=0.97, IFI)=0.99, CFI=0.99, RMSEA=0.03, AGFI=0.93	Cronbach's alpha: 0.93 to 0.95, CR=0.91 to 0.94	115	Physicians	Hospital
17	Carayon et al. (Model of electronic health records (EHR) acceptance) (25)	122	MRA	-	Cronbach's alpha: 0.90-0.98	282	Nurses	Medical Center
19	Tilahun et al. (26)	109	CFA-SEM	AVE=0.68 to 0.81, CR=0.84 to 0.96, $\chi^2/d.f. = 2.39$, GFI = 0.92, AGFI = 0.87, NFI = 0.92, RMSR = 0.056	Cronbach's alpha:0.84 to 0.91	384	Physicians, nurses, and other clinicians	Hospital
20	Steininger et al. (27)	506	SEM	$\chi^2 = 38.76$, df=13, AIC=100.7, CFI=60.97, TLI=0.91, RMSEA=0.10	Cronbach's alpha: 0.76 to 0.92	204	Physicians	Physician Office
21	Kowitzlawakulet al. (28)	32	SEM	$\chi^2/d.f.=1.757$, GFI=0.90, NFI=0.97, IFI=0.931, CFI=0.969, RMSEA=0.06, AGFI=0.863	Cronbach's alpha: 0.76 to 0.92	212	Nurses	Primary Care Clinic
22	Hsieh et al. (29)	511	CFA	AVE=0.766 to 0.961, CR=0.908 to 0.967	not reported	191	Physicians	Hospital
23	Abdekhoda et al. (30)	537	SEM	Relative $\chi^2 = 1.9$, df=13, NFI =0.93, CFI=0.93, TLI=0.92, RMSEA=0.02	not reported	234	Physicians	Hospital
24	Messeri et al. (31)	7	MRA	-	Cronbach's alpha: 0.64 to 0.74	460	Physicians and nurses	Primary Care Clinic
25	Hsu et al. (32)	56	MRA	-	Cronbach's alpha: 0.622 to 0.930, CR: 0.650 to 0.930	720	Nurses	Hospital
26	Leblanc et al. (33)	53	MRA	-	Cronbach's alpha: 0.62 to 0.89	99	Nurses	Hospital
27	Chen and Hsiao (34)	50	CFA, MRA	AVE: > 0.51	CR: > 0.87	81	Physicians	Hospital
28	Hsiao et al. (35)	26	MRA	-	Cronbach's alpha: 0.97	545	Nurses	Hospital
29	Sicotte et al. (36)	33	Linear regression analysis	-	Cronbach's alpha: 0.73 and 0.97	125	Physicians	Hospital
30	Erlirianto et al. (37)	602	SEM	FIT: 0.386, AFIT=0.370, GFI = 0.943	not reported	67	Nurses, other clinicians, and other non-clinical participants	Hospital
31	Ho et al. (38)	11	Pearson correlations	-	Cronbach's alpha: > 0.87	581	Physicians	Primary Care Clinic
32	Devine et al. (39)	52	Linear regression analysis	-	Cronbach's alpha: 0.90 to 0.92	117	Physicians and nurses	Primary Care Clinic
33	Otieno et al. (composite index (CI)) (40)	64	Correlation analysis	-	Cronbach's alpha: 0.475 to 0.843	1892	Physicians, nurses, and other non-clinical participants	Hospital
34	Otieno et al. (41)	66	EFA	KMO >0.80 (Field 2005) and Bartlett's tests P-value >0.00	Cronbach's alpha total:0.88, range: 0.36 to 0.94	1666	Physicians, nurses, and other non-clinical participants	Hospital

EFA
correlation analysis CA
Multiple Regression Analysis MRA

Data gathering methods, missing data, and outlier handling

Data were collected using various methods, including in-person (n=14), online (n=11), in-person or online (n=4), email (n=3), and nuclear (n=6). Handling missing cases was discussed in 20.6% of papers (n=7). Therefore, only one of the selected papers explained outlier handling.

Analysis software

The most commonly used software was SPSS, used in 14.7% of the papers. AMOS and Smart PLS were used in 11.8% and 8.8% of the papers, respectively, followed by GeSCA, LISREL, SAS, and STAT, each used in less than 6% of papers. It is worth noting that used software not reported was used in 35.3% (n=12) of papers.

Theoretical Basis models

A number of 16 studies (47%) used the Technology Acceptance Model (TAM) to develop the EHR evaluation model, while the Information System Success Model (ISSM) was used in 7 papers (20%). Other models used to develop the EHR evaluation model included the Ammenwerth and Dekeizer (ADK) model, Computer Anxiety Model (CAM), Diffusion of Innovation (DOI), Composite Index (CI), Dual Factor Model (DFM), Fit Between Individuals, Task, and Technology framework (FITT), Human, Organization, and Technology-Fit Factors (HOT-Fit), Job Demands Resource Model (JDRM), Theory of Planned Behavior (TPB), Task and Technology and Fit (TTF), Technology Readiness Index (TRI), as well as Unified Theory of Acceptance and Use of Technology (UTAUT).

Relationship between variables of the EHR evaluation models

Among 34 papers, 29 articles (85.29%) developed a causal model that included independent, moderating, and dependent variables. Four causal models did not include moderating variables, while five papers proposed a non-causal model for evaluating Electronic Health Records (EHR). The EHR evaluation model diagram was included in most articles (n=27). The moderating variables that were most commonly observed in the analyzed articles were perceived usefulness (n=12) and perceived ease of use (n=9). The dependent variables that received the most attention were behavior intention to use/system use (n=16) and system adoption (n=6). The independent variables that were most frequently studied were system quality (n=11), information quality (n=7), and service quality (n=6).

Statistical validation methods

Among the 34 papers that disclosed the validation methods utilized, the most widely used technique was SEM, which was employed in 26.5% (n=9) of the studies. Other methods utilized were Confirmatory Factor Analysis (CFA) (17.6%; n=6), Exploratory Factor Analysis (EFA) (11.8%; n=4), Multiple Regression Analysis (MRA) (14.7%; n=5), Correlation (5.9%; n=2), Linear Regression (5.9%; n=2), and Pearson (2.9%; n=1). Furthermore, it is worth noting that in addition to the methods listed in the articles, four studies used both CFA-SEM (11.8%) and CFA, while one study (2.9%) utilized both CFA and MRA.

Sampling technique and sample size

Convenience sampling was the most frequently used method of sampling, reported in 73.5% of papers, followed by random sampling (5.9%) and bootstrapping resampling (2.9%). Six papers (17.6%) did not provide information on the sampling technique used. The mean sample size used in the papers was 504.14 (SD= ±570.99, range = 51 to 2590). Among the 34 papers, the majority (n=25; 73.5%) used convenience sampling as their sampling technique, which was the most frequently used method. Random sampling was used in only 2 (5.9%) papers, while bootstrapping resampling was used in only 1 (2.9%) paper. Meanwhile, 6 (17.6%) papers did not provide information on the sampling technique used. The means and 95% confidence intervals for each of the different validation methods were in ascending order as follows: Linear regression: Mean = 121(SD=±4), CFA and SEM: Mean = 249.5 (SD=±58.23), CFA: Mean = 305.33 (SD=±121.82), EFA: Mean = 662.75 (SD=±336.52), MRA: Mean = 421.2 (SD=±107.7), SEM: Mean = 629 (SD=±267.62), and Correlation: Mean = 1494.5, (SD=±397.5).

Reliability assessment methods and values

A reliability assessment was performed in 28 (82%) articles. Cronbach's alpha and composite reliability (CR) were popular reliability (internal

validation) methods on identified papers. Reliability was not reported in six articles. Cronbach's alpha and composite reliability (CR) were calculated together in seven articles, while Cronbach's alpha was calculated in 25 (73.5%) papers. The CR was calculated in 9 (26%) articles. The mean of CR was 0.94 (SD=±0.03), with values ranging from 0.87-1.00 in identified studies. As for Cronbach's alpha, the mean was 0.92 (SD=±0.058), varying from 0.74-1.00.

Statistics criteria in the SEM approach

The most commonly used criteria in the SEM approach included the Comparative Fit Index (CFI), Root Mean Square Error of Approximation (RMSEA), Chi-square (χ^2), and Normed Fit Index (NFI). After analyzing eleven articles, it was found that the mean CFI was 6.30845, with a range of 0.015-60.97 and a standard deviation of 18.13. The mean RMSEA was 0.172 (SD=±0.29) across nine papers. The Chi-square (χ^2) was reported in eight articles, with values ranging from 1.90-3852.620. The NFI in seven papers was 0.94 (SD=±0.3). The Degree of Freedom (df) was calculated in six articles, ranging from 1.82-466.00. The Goodness-of-Fit Index (GFI) had a mean of 0.85 (SD=±0.14) across five papers. The χ^2/df was reported in four articles, with values ranging from -2.72-2.41. The TLI was measured in five articles, with values ranging from 0.91-0.93 and a mean of 0.918 (SD=±0.008). The Incremental Fit Index (IFI) in four papers ranged from 0.931-0.990.

The Non-Normed Fit Index (NNFI) was calculated in two papers, with values of 0.91 and 0.98. Differences in CFI were reported in one article, with a value of 0.9900. The range of the Adjusted Goodness-of-Fit Index (AGFI) in three papers was 0.86-0.93. The Root Mean Square Residual (RMR) was reported in three papers, ranging from 0.035-0.928. The Standardized Root Mean Square Residual (SRMR) ranged from 0.048-0.056 in two articles. The value of the Parsimony Goodness-of-Fit Index (PGFI) in one paper was 0.68. The Akaike Information Criterion (AIC) was reported in one article, with a value of 100.700. None of the identified studies reported criteria, such as PCFI, PNFI, RFI, P Ratio, NCP, FIMIN, RMSEA with a confidence interval, PCLOSE, CAIC, BCC, BIC, ECVI, and MECVI.

The most commonly used criteria in the SEM approach included the Comparative Fit Index (CFI), Root Mean Square Error of Approximation (RMSEA), Chi-square (χ^2), and Normed Fit Index (NFI). After analyzing eleven articles, it was revealed that the mean CFI was 6.30845, with a range of 0.015-60.97 and a standard deviation of 18.13. The mean RMSEA was 0.172 (SD=±0.29) across nine papers. The Chi-square (χ^2) was reported in eight articles, with values ranging from 1.90-3852.620. The NFI in seven papers was 0.94 (SD=±0.3). The Degree of Freedom (df) was calculated in six articles, ranging from 1.82-466.00. The Goodness-of-Fit Index (GFI) had a mean

of 0.85 (SD=±0.14) across five papers. The χ^2/df was reported in four articles, with values ranging from 2.72-2.41. TLI was measured in five articles, with values ranging from 0.91-0.93 and a mean of 0.918 (SD=±0.008). The four papers, Incremental Fit Index (IFI) ranged from 0.931-0.990.

The Non-Normed Fit Index (NNFI) was calculated in two papers, with values of 0.91 and 0.98. Differences in CFI were reported in one article, with a value of 0.9900. The range of the Adjusted Goodness-of-Fit Index (AGFI) in three papers was 0.86-0.93. The Root Mean Square Residual (RMR) was reported in three papers, ranging from 0.035-0.928. The Standardized Root Mean Square Residual (SRMR) ranged from 0.048-0.056 in two articles. The Parsimony Goodness-of-Fit Index (PGFI) value in one paper was 0.68. The Akaike Information Criterion (AIC) was reported in one article, with a value of 100.700.

None of the identified studies reported such criteria as PCFI, PNFI, RFI, P Ratio, NCP, FIMIN, RMSEA with a confidence interval, PCLOSE, CAIC, BCC, BIC, ECVI, and MECVI. A reliability assessment was performed in 28 (82%) articles. Cronbach's alpha and composite reliability (CR) were popular reliability (internal validation) methods on identified papers. Reliability was not reported in six articles. Cronbach's alpha and composite reliability (CR) were calculated together in seven articles, while Cronbach's alpha was calculated in 25 (73.5%) articles. The CR was calculated in 9 (26%) articles. The mean of CR was 0.94 (SD=±0.03), with values ranging from 0.87-1.00 in identified studies. As for Cronbach's alpha, the mean was 0.92 (SD=±0.058), varying from 0.74-1.00. A reliability assessment was performed in 28 (82%) articles. Cronbach's alpha and composite reliability (CR) were popular reliability (internal validation) methods on identified papers. Reliability was not reported in six articles.

Cronbach's alpha and composite reliability (CR) were calculated together in seven articles, while Cronbach's alpha was calculated in 25 (73.5%) articles. The CR was calculated in 9 (26%) articles. The mean of CR was 0.94 (SD=±0.03), with values ranging from 0.87-1.00 in identified studies. As for Cronbach's alpha, the mean was 0.92 (SD=±0.058), varying from 0.74-1.00. The most commonly used criteria in the SEM approach included the Comparative Fit Index (CFI), Root Mean Square Error of Approximation (RMSEA), Chi-square (χ^2), and Normed Fit Index (NFI). After analyzing 11 articles, it was found that the mean CFI was 6.30845, with a range of 0.015-60.97 and a standard deviation of 18.13. The mean RMSEA was 0.172 (SD=±0.29) across nine papers. The Chi-square (χ^2) was reported in eight articles, with values ranging from 1.90-3852.620. The NFI in seven papers was 0.94 (SD=±0.3).

The Degree of Freedom (df) was calculated in six articles, ranging from 1.82-466.00. The Goodness-of-Fit Index (GFI) had a mean of 0.85 (SD=±0.14) across

five papers. The χ^2/df was reported in four articles, with values ranging from -2.72-2.41. TLI was measured in five articles, with values ranging from 0.91-0.93 and a mean of 0.918 (SD=±0.008). The Incremental Fit Index (IFI) in four papers ranged from 0.931-0.990.

The Non-Normed Fit Index (NNFI) was calculated in two papers, with values of 0.91 and 0.98. Differences in CFI were reported in one article, with a value of 0.9900. The range of Adjusted Goodness-of-Fit Index (AGFI) in three papers was 0.86 to 0.93. The Root Mean Square Residual (RMR) was reported in three papers, ranging from 0.035-0.928. The Standardized Root Mean Square Residual (SRMR) ranged from 0.048-0.056 in two articles. The Parsimony Goodness-of-Fit Index (PGFI) value in one paper was 0.68. The Akaike Information Criterion (AIC) was reported in one article, with a value of 100.700. None of the identified studies reported criteria, such as PCFI, PNFI, RFI, P Ratio, NCP, FIMIN, RMSEA with a confidence interval, PCLOSE, CAIC, BCC, BIC, ECVI, and MECVI.

Statistics criteria in CFA and EFA approach

The AVE in 10 papers was calculated. The articles analyzed had a mean of 0.71 (SD=±0.97) for lower AVE with a minimum value of 0.51 and a maximum value of 0.81. The mean for upper AVE was 0.8863 (SD=±0.97), with a range of 0.80-1.00. KMO was measured in three papers. Bartlett's test was significant in all these studies.

5. Discussion

The present study aimed to conduct an overview of reliability and validity evaluation methods in EHR evaluation models. The obtained results revealed that EHR evaluation models were created using various validation methods, each with different levels of accuracy. In addition, the effectiveness of these methods depended on the sample size used in the study. Researchers can perform more reliable validation methods in studies with larger sample sizes. Our research revealed that the most common validation methods applied in EHR evaluation models are SEM and CFA. Factor analysis and structural analysis are the two fundamental methods used for evaluating models, with high power to identify the model structure and variable relationships (7). The SEM is a crucial tool for analyzing and comprehending causal relationships among direct and indirect variables in EHR evaluation studies (6, 42). It has been applied to evaluate EHR implementation and its impact on users in previous studies (43). The SEM can provide a new perspective on analyzing data and the potential for advancing research in medical and health sciences (44). In agreement with our findings, Dash and Paul pointed out that SEM can overcome the limitations of

mediation analysis, CFA, and regression methods (5) and has widespread use in social and human studies (45). The SEM has been used in various healthcare studies, such as those on COVID-19 (46), nursing research (47) and psychological research (48).

SEM offers a new perspective on data analysis and has the potential to advance research in medical and health sciences. (44). Dash and Paul pinpointed that SEM can overcome the limitations of mediation analysis, CFA, and regression methods (5), and it is widely used in social and human studies (45). SEM has also been applied in various healthcare studies, including research on COVID-19 (46), augmented reality applications (46), nursing research (47), and psychological research (48). Based on our findings, it can be observed that numerous studies have developed causal models using independent, moderating, and dependent variables, leading to the widespread use of SEM as a validation method. This is further supported by previous research, such as the study by Ismael and Duleba, who suggested that SEM allows for the measurement of observed variables during data collection while indirect connections can be made to examine latent variables (49).

Based on the results of the current study, a number of identified studies used both CFA and SEM as validation methods. These findings are supported by previous research, such as the model proposed by Dash and Paul, which outlines a five-step process for utilizing SEM. This process includes individual construct identification, preparation for CFA, running CFA, structural modeling, and finally, interpreting findings (5). In a similar vein, Dragan and Topolsek have outlined the basic steps often used in SEM modeling, which include EFA, CFA, and calculating model indices (50). The CFA is a crucial component of the measurement model, which assesses the relationships between a construct and its indicator variables (7). This process helps explain how the data fits into the structure model (11). Once the measurement model is validated using CFA, pathway analysis is conducted to examine the relationships among the latent variables (5). Path modeling involves estimating multiple regression models simultaneously and can demonstrate mediation, moderation, and interaction relationships among variables (5).

Our review revealed that some of the identified studies used EFA to develop an EHR validation model. In the study by Dragan and Topolsek, EFA was used as a preliminary step in structural modeling when the researchers were interested in data reduction. It is essential to note that the selection of analysis methods may vary depending on the research question and goals of the study (50). Furthermore, our findings demonstrated that the goodness of fit index (GFI), adjusted goodness of fit index (AGFI), and root mean square error of approximation (RMSEA) criteria were prominent in all the included

criteria for SEM approaches. This is in line with previous studies, where GFI, AGFI, and RMSEA measures were commonly used in the SEM literature to assess the goodness of fit of the model (48).

TAM, TPB, and UTAUT are all human and social-based theories that focus on understanding individuals' attitudes toward technology adoption (51). These theories were used frequently in our identified papers. In healthcare research, Technology Acceptance Model (TAM), as well as its extensions and modifications, are at the forefront. Modified and extended TAM models were developed to improve the explanatory power of the original TAM model. In addition, the Unified Theory of Acceptance and Use of Technology (UTAUT) is considered one of the other most pertinent and actively used models in technology acceptance studies in healthcare (52). The obtained results pointed to several ways to develop an EHR evaluation framework. A framework can be developed based on a literature review or a combination of existing frameworks. For instance, measuring end-user computing satisfaction (EUCS) was developed based on UTAUT, Doll and Torkzadeh (1988), and Bailey and Pearson's model (1983) (22). End-User Acceptance Model (EUAM) is a hybrid framework of TAM and UTAUT (10). Lambooi et al. carried out a literature review and developed an original evaluation framework, which was not based on any existing evaluation framework (8). Michel-Verkerke et al. combined observations of the developments in the EHR market and healthcare together with a literature search to propose the "Six P-model of EPR-use" (16).

Another method to develop an EHR evaluation framework is to modify an existing EHR evaluation framework which refers to adding or removing dimensions to an existing evaluation framework. For example, Sicotte et al. added two variables to ISSM that had been used in another study on the benefits of PACS (36). Messeri et al. conducted a literature search and prepared a modified ISSM (31). Chen and Hsiao extended TAM to take quality aspects into account (34). Tilahun and Fritz added computer literacy to ISSM (26). Steininger and Stiglbauer presented a modified TAM and included social influence, health information technology experience, and privacy concern factors as external variables to the TAM [27].

6. Conclusion

It is believed that the results of this study can assist researchers in examining and modifying EHR evaluation frameworks to suit their specific needs. Furthermore, our findings serve as a solid foundation for the development of new EHR evaluation frameworks. The present review study can assist researchers in understanding the viability of validation methods, such as SEM techniques,

correlation models, and hierarchical regression models in the context of electronic health records. Moreover, it is recommended that researchers utilize the findings presented in this article to enhance the implementation and utilization of SEM, CFA, and EFA methods in EHR evaluation models.

Acknowledgments

None.

Conflicts of interest

The authors declare that they have no conflict of interest.

References

- Nykänen P, Brender J, Talmon J, de Keizer N, Rigby M, Beuscart-Zephir MC, Ammenwerth E. Guideline for good evaluation practice in health informatics (GEP-HI). *Int J Med Inform.* 2011;**80**(12):815-27. doi: [10.1016/j.ijmedinf.2011.08.004](https://doi.org/10.1016/j.ijmedinf.2011.08.004). [PubMed: 21920809].
- Sokolow PS, Crawford PR, Lehmann HP. Health services research evaluation principles. *Methods Inf Med.* 2012;**51**(02):122-30. doi: [10.3414/me10-01-0066](https://doi.org/10.3414/me10-01-0066). [PubMed: 22311125].
- Rahimi B, Nadri H, Afshar HL, Timpka T. A systematic review of the technology acceptance model in health informatics. *Appl Clin Inform.* 2018;**9**(03):604-34. doi: [10.1055/s-0038-1668091](https://doi.org/10.1055/s-0038-1668091). [PubMed: 30112741].
- Ebnehoseini Z, Tabesh H, Jangi MJ, Deldar K, Mostafavi SM, Tara M. Investigating evaluation frameworks for electronic health record: a literature review. *Open Access Maced J Med Sci.* 2021;**9**(E):8-25. doi: [10.3889/oamjms.2021.3421](https://doi.org/10.3889/oamjms.2021.3421).
- Dash G, Paul J. CB-SEM vs PLS-SEM methods for research in social sciences and technology forecasting. *Technol Forecast Soc Change.* 2021;**173**:121092. doi: [10.1016/j.techfore.2021.121092](https://doi.org/10.1016/j.techfore.2021.121092).
- Deng L, M. Yang and K.M. Marcoulides, Structural equation modeling with many variables: A systematic review of issues and developments. *Front Psychol.* 2018;**9**:580. doi: [10.3389/fpsyg.2018.00580](https://doi.org/10.3389/fpsyg.2018.00580). [PubMed: 29755388].
- Ghanbar H, Rezvani R. Structural Equation Modeling in L2 Research: A Systematic Review. *International Journal of Language Testing.* 2023 Mar 1;**13**(Special Issue):79-108. doi: [10.22034/ijlt.2023.381619.1224](https://doi.org/10.22034/ijlt.2023.381619.1224).
- Lambooi, M.S., H.W. Drewes and F. Koster, Use of electronic medical records and quality of patient data: different reaction patterns of doctors and nurses to the hospital organization. *BMC Med Inform Decis Mak.* 2017;**17**(1):1-11. doi: [10.1186/s12911-017-0412-x](https://doi.org/10.1186/s12911-017-0412-x). [PubMed: 28187729].
- Mohd Salleh, M.I., N. Zakaria and R. Abdullah, The influence of system quality characteristics on health care providers' performance: Empirical evidence from Malaysia. *Journal of infection and public health.* 2016;**9**(6): 698-707. doi: [10.1016/j.jiph.2016.09.002](https://doi.org/10.1016/j.jiph.2016.09.002). [PubMed: 27659115].
- Kim, S., K.H. Lee, H. Hwang and S. Yoo, Analysis of the factors influencing healthcare professionals' adoption of mobile electronic medical record (EMR) using the unified theory of acceptance and use of technology (UTAUT) in a tertiary hospital. *Med Inform Decis Mak.* 2016;**16**(1). doi: [10.1186/s129117](https://doi.org/10.1186/s129117). [PubMed: 26831123].
- Liu, C.F. and T.J. Cheng, Exploring critical factors influencing physicians' acceptance of mobile electronic medical records based on the dual-factor model: a validation in Taiwan. *BMC Med Inform Decis Mak.* 2015. **15**:4. doi: [10.1186/s12911-014-0125-3](https://doi.org/10.1186/s12911-014-0125-3). [PubMed: 25889506].
- Sintonen, S., K. Mäkelä and R. Miettinen, User acceptance of electronic health records: A post-implementation study. *International Journal of Healthcare Technology and Management.* 2015;**15**(2):162-75. doi: [10.1504/IJHTM.2015.074556](https://doi.org/10.1504/IJHTM.2015.074556).
- Gan, Q., Is the adoption of electronic health record system "contagious"? *Health Policy Technol.* 2015. **4**(2): 107-112. doi: [10.1016/j.hlpt.2015.02.009](https://doi.org/10.1016/j.hlpt.2015.02.009).
- Kralj, D., J. Kern, S. Tonkovic and M. Koncar, Development of the quality assessment model of EHR software in family medicine practices: research based on user satisfaction. *J Innov Health Inform.* 2015;**22**(3): 340-58. doi: [10.14236/jhi.v22i3.158](https://doi.org/10.14236/jhi.v22i3.158). [PubMed: 26577425].
- Wang, J.Y., H.Y. Ho, J.D. Chen, S. Chai, C.J. Tai and Y.F. Chen, Attitudes toward inter-hospital electronic patient record exchange: discrepancies among physicians, medical record staff, and patients. *BMC Health Serv Res.* 2015. **15**: p. 264. doi: [10.1186/s12913-015-0896-y](https://doi.org/10.1186/s12913-015-0896-y). [PubMed: 26162748].
- Michel-Verkerke, M.B., R.A. Stegwee and T.A.M. Spil, The six P's of the next step in electronic patient records in the Netherlands. *Health Policy Technol.* 2015;**4**(2): 137-43. doi: [10.1016/j.hlpt.2015.02.011](https://doi.org/10.1016/j.hlpt.2015.02.011).
- Hysong, S.J., C. Spitzmuller, D. Espadas, D.F. Sittig and H. Singh, Electronic alerts and clinician turnover: the influence of user acceptance. *Am J Manag Care.* 2014;**20**(17): p. SP520-SP530. [PubMed: 25811826].
- Gagnon, M.P., K. Ghandour el, P.K. Talla, D. Simonyan, G. Godin, M. Labrecque, et al., Electronic health record acceptance by physicians: testing an integrated theoretical model. *J Biomed Inform.* 2014;**48**:17-27. doi: [10.1016/j.jbi.2013.10.010](https://doi.org/10.1016/j.jbi.2013.10.010). [PubMed: 24184678].
- Iqbal, U., C.H. Ho, Y.C. Li, P.A. Nguyen, W.S. Jian and H.C. Wen, The relationship between usage intention and adoption of electronic health records at primary care clinics. *Computer Methods and Programs in Biomedicine.* 2013;**112**(3): 731-737. doi: [10.1016/j.cmpb.2013.09.001](https://doi.org/10.1016/j.cmpb.2013.09.001). [PubMed: 24091088].
- Kuo, K.M., C.F. Liu and C.C. Ma, An investigation of the effect of nurses' technology readiness on the acceptance of mobile electronic medical record systems. *Med Inform Decis Mak.* 2013;**13**(1). doi: [10.1186/1472-6947-13-88](https://doi.org/10.1186/1472-6947-13-88). [PubMed: 23938040].
- Lu, C.H., J.L. Hsiao and R.F. Chen, Factors determining nurse acceptance of hospital information systems. *Comput Inform Nurs.* 2012;**30**(5): 257-64. doi: [10.1097/NCN.0b013e318224b4cf](https://doi.org/10.1097/NCN.0b013e318224b4cf). [PubMed: 22228251].
- Aggelidis, V.P. and P.D. Chatzoglou, Hospital information systems: measuring end user computing satisfaction (EUCS). *J Biomed Inform.* 2012;**45**(3): 566-79. doi: [10.1016/j.jbi.2012.02.009](https://doi.org/10.1016/j.jbi.2012.02.009). [PubMed: 22426283].
- Chen, R.F. and J.L. Hsiao, An investigation on physicians' acceptance of hospital information systems: a case study. *Int J Med Inform.* 2012;**81**(12): 810-20. doi: [10.1016/j.ijmedinf.2012.05.003](https://doi.org/10.1016/j.ijmedinf.2012.05.003). [PubMed: 22652011].
- Lin, C., I.C. Lin and J. Roan, Barriers to physicians' adoption of healthcare information technology: An empirical study on multiple hospitals. *J Med Syst.* 2012;**36**(3): 1965-77. doi: [10.1007/s10916-011-9656-7](https://doi.org/10.1007/s10916-011-9656-7). [PubMed: 21336605].
- Carayon, P., R. Cartmill, M.A. Blosky, R. Brown, M. Hackenberg, P. Hoonakker, et al., ICU nurses' acceptance of electronic health records. *Journal of the American Medical Informatics Association : JAMIA.* 2011;**18**(6): 812-9. doi: [10.1136/amiajnl-2010-000018](https://doi.org/10.1136/amiajnl-2010-000018). [PubMed: 21697291].
- Tilahun, B. and F. Fritz, Modeling antecedents of electronic medical record system implementation success in low-resource setting hospitals. *Med Inform Decis Mak.* 2015;**15**: 61. doi: [10.1186/s12911-015-0192-0](https://doi.org/10.1186/s12911-015-0192-0). [PubMed: 26231051].
- Steininger, K. and B. Stiglbauer, EHR acceptance among Austrian resident doctors. *Health Policy Technol.* 2015;**4**(2): p. 121-130. doi: [10.1016/j.hlpt.2015.02.003](https://doi.org/10.1016/j.hlpt.2015.02.003).
- Kowitlawakul Y, S.W. Chan, J. Pulcini, W Wang, Factors influencing nursing students' acceptance of electronic health records for nursing education (EHRNE) software program. *Nurse education today.* 2015;**35**(1): 189-94. doi: [10.1016/j.nedt.2014.05.010](https://doi.org/10.1016/j.nedt.2014.05.010). [PubMed: 24947068].
- Hsieh, P.J., Physicians' acceptance of electronic medical records exchange: An extension of the decomposed TPB model with

- institutional trust and perceived risk. *Int J Med Inform*, 2015;**84**(1): 1-14. doi:10.1016/j.ijmedinf.2014.08.008. [PubMed:25242228].
30. Abdekhoda, M., M. Ahmadi, M. Gohari and A. Noruzi, The effects of organizational contextual factors on physicians' attitude toward adoption of Electronic Medical Records. *J Biomed Inform*, 2015, 53: p. 174-179. doi:10.1016/j.jbi.2014.10.008. [PubMed: 25445481].
 31. Messeri, P., S. Khan, M. Millery, A. Campbell, J. Merrill, S. Shih, et al., An information systems model of the determinants of electronic health record use. *Appl Clin Inform*, 2013; **4**(2): p. 185-200. doi:10.4338/aci-2013-01-ra-0005. [PubMed:23874357].
 32. Hsu, S.C., C.F. Liu, R.H. Weng and C.J. Chen, Factors influencing nurses' intentions toward the use of mobile electronic medical records. *Comput Inform Nurs*, 2013;**31**(3): 124-32. doi:10.1097/NXN.0b013e318270100b. [PubMed:23114391].
 33. Leblanc, G., M.P. Gagnon and D. Sanderson, Determinants of primary care nurses' intention to adopt an electronic health record in their clinical practice. *Comput Inform Nurs*, 2012;**30**(9): 496-502. doi:10.1097/NXN.0b013e318257db17. [PubMed:22592453].
 34. Chen, R.F. and J.L. Hsiao, An empirical study of physicians' acceptance of hospital information systems in Taiwan. *Telemedicine journal and e-health : Telemed J E Health*, 2012;**18**(2): 120-5. doi:10.1089/tmj.2011.0081. [PubMed:22283362].
 35. Hsiao, J.L., H.C. Chang and R.F. Chen, A study of factors affecting acceptance of hospital information systems: a nursing perspective. *J Nurs Res*. 2011;**19**(2): p. 150-60. doi:10.1097/JNR.0b013e31821cbb25. [PubMed:21586992].
 36. Sicotte, C., G. Pare, K.K. Bini, M.P. Moreault and G. Laverdure, Virtual organization of hospital medical imaging: a user satisfaction survey. *J Digit Imaging*, 2010;**23**(6): 689-700. doi:10.1007/s10278-009-9220-x. [PubMed:19588196].
 37. Erlirianto LM, AHN Ali , A Herdiyanti. The Implementation of the Human, Organization, and Technology-Fit (HOT-Fit) Framework to Evaluate the Electronic Medical Record (EMR) System in a Hospital. *Procedia Comput Sci*, 2015;**72**: 580-7. doi:10.1016/j.procs.2015.12.166.
 38. Ho, C.H., H.C. Wene, C.M. Chu, Y.S. Wu and J.L. Wang, Importance-satisfaction analysis for primary care physicians' perspective on EHRs in Taiwan. *Int J Environ Res Public Health*, 2014;**11**(6): 6037-51. doi:10.3390/ijerph110606037. [PubMed:24914640].
 39. Devine, E.B., R. Patel, D.R. Dixon and S.D. Sullivan, Assessing attitudes toward electronic prescribing adoption in primary care: a survey of prescribers and staff. *Inform Prim Care*, 2010;**18**(3): 177-87. [PubMed:21396241].
 40. Otieno GO, T Hinako, Motohiro K, Daisuke , N Keiko. Measuring effectiveness of electronic medical records systems: towards building a composite index for benchmarking hospitals. *Int J Med Inform*, 2008;**77**(10): 657-69. doi:10.1016/j.ijmedinf.2008.01.002. [PubMed: 18313352].
 41. Otieno, O.G., H. Toyama, M. Asonuma, M. Kanai-Pak and K. Naitoh, Nurses' views on the use, quality and user satisfaction with electronic medical records: questionnaire development. *J Adv Nurs*, 2007;**60**(2): 209-19. doi:10.1111/j.1365-2648.2007.04384.x. [PubMed:17877568].
 42. Cho KW, SK Bae, JH. Ryu, KN Kim, CH An ,YM Chae, Performance evaluation of public hospital information systems by the information system success model. *Healthc Inform Res*, 2015;**21**(1): 43-8. doi:10.4258/hir.2015.21.1.43. [PubMed:25705557].
 43. Bahadori MK, Alimohammadzadeh K. Abdolkarimi, RRavangard, Factors affecting physicians' attitudes towards the implementation of electronic health records using structural equation modeling modeling (SEM). *Shiraz E Medical Journal*, 2017; **18**(11). doi: 10.5812/semj.13729.
 44. Beran, T.N. and C. Violato, Structural equation modeling in medical research: a primer. *BMC Res Notes*, 2010; **3**(1): 1-10. doi: 10.1186/1756-0500-3-267. [PubMed:20969789].
 45. Shaheen, F., N. Ahmad, M. Waqas, A. Waheed and O. Farooq, Structural equation modeling (SEM) in social sciences & medical research: a guide for improved analysis. *Int. J Acad Res Bus Soc Sci*. 2017;**7**(5): 132-43. doi: 10.6007/IJARBS/v7-i5/2882.
 46. Nguyen, V.T. and C.T.H. Nguyen, A systematic review of structural equation modeling in augmented reality applications. *arXiv preprint arXiv:2301.11811*, 2023. doi:10.48550/arXiv.2301.11811.
 47. Sharif, S.P., I. Mostafiz and V. Guptan, A systematic review of structural equation modelling in nursing research. *Nurse Res*, 2019;**30**(2). doi: 10.7748/nr.2018.e1577. [PubMed:30207432].
 48. MacCallum, RC., JT. Austin. Applications of structural equation modeling in psychological research. *Annu Rev Psychol*, 2000;**51**(1): 201-26. doi:10.1146/annurev.psych.51.1.201. [PubMed:10751970].
 49. Ismael, K. and S. Duleba, A Systematic Review of the Latest Advancements on Structural Equation Modelling (SEM) Technique Focusing on Applications in Transportation Planning. *Period. Polytech. Transp. Eng*, 2022;**50**(4): 336-43. doi:10.3311/PPtr.17523.
 50. Dragan D, D Topolšek. Introduction to structural equation modeling: review, methodology and practical applications. *The International Conference on Logistics & Sustainable Transport*. 2014.
 51. Venkatesh V, MG Morris, GB Davis , FD Davis. User acceptance of information technology: Toward a unified view. *MIS quarterly*. 2003: 425-78. doi: 10.2307/30036540.
 52. Kruse CS, K Kothman, K Anerobi, L Abanaka. Adoption Factors of the Electronic Health Record: A Systematic Review. *JMIR medical informatics*, 2016;**4**(2): e19. doi:10.2196/medinform.5525. [PubMed:27251559].