Published online 2025 April

Effect of Emotion Regulation Training Based on Gross Model on Psychological Well-being and Sleep Quality in Women with Psychosomatic Disorders (Rheumatic and Joint Pain)

Maryam Dastjani Farahani¹, Azadeh Tavoli^{1*}, Elahe Mirzayi Sefidabi²

1. Department of Psychology, Faculty of Education and Psychology, Alzahra University, Tehran, Iran

2. Department of Psychology, Payame Noor University (PNU), Tehran, Iran

* **Corresponding author:** Azadeh Tavoli, Department of Psychology, Faculty of Education and Psychology, Alzahra University, Tehran, Iran. Email: a.tavoli@alzahra.ac.ir

Received:26 March 2024 Accepted: 17 December 2024

Abstract

Background: Rheumatoid arthritis is a chronic, systemic, inflammatory disease classically affecting the small joints of the hands and feet.

Objectives: The purpose of this research was to investigate the effect of emotion regulation training based on the Gross model on the psychological well-being and sleep quality of women with psychosomatic disorders (rheumatic and joint pains).

Methods: This semi-experimental study was conducted based on a pretest-posttest design and a control group. The statistical population of this study consisted of all women with psychophysical disorders in Farahan City, among which 40 individuals were selected through the convenience sampling method and then randomly allocated to either the experimental or control group (n=20 each). The required data were collected using the Ryff Psychological Well-Being and Pittsburgh Sleep Quality Index. Emotion regulation techniques were taught using Gross's standard training protocol in eight 90-minute sessions, while the control group was on the waiting list. The data were analyzed using the analysis of covariance in SPSS-23 software.

Results: The study's results indicate that instructing women with psychosomatic disorders on how to handle their emotions using the Gross model can greatly enhance their mental health (P<0.01). Moreover, the findings from the analysis of covariance revealed that Emotion Regulation Training based on the Gross model led to enhancements in sleep quality in women with psychosomatic disorders (P<0.01).

Conclusion: The outcomes of this study have the potential to enhance the overall well-being and sleep quality of individuals experiencing rheumatic and joint pains.

Keywords: Emotion regulation training, Gross model, Psychological well-being, Sleep quality

1. Background

Psychosomatic disorders are a collection of disorders that develop due to mental factors; essentially, they manifest in the body due to mental strain, stress, and chronic pressure (1). These disorders vary in presentation from person to person of any age or circumstance; however, they are most commonly seen in individuals aged 20 to 45 years. Experiencing various mental stresses and issues can lead these individuals to exhibit psychosomatic symptoms and experience uncomfortable bodily pains known as psychosomatic pains (2). Among the psychosomatic pains prevalent today are rheumatic and joint pain. The onset of rheumatic and joint pain typically involves episodes of joint inflammation, leading to swelling and discomfort. This disease causes gradual harm to the musculoskeletal system

Copyright © 2025, Razavi International Journal of Medicine. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

and results in pain and permanent damage to bone and cartilage, which elicits various responses. Pain experienced by these patients can lead to emotions of inadequacy, worry, anxiety, and stress, especially when it continues for an extended period (3). As stated in a study by Aslund et al. (2014), one of the mental factors affected by chronic pain in individuals is their psychological well-being (4). Mental well-being refers to the level of and positive functioning and personal represents the dimensions of assessment regarding satisfaction with life (5).

According to a study, individuals with chronic pain experience a decline in their psychological well-being as a result of the psychological stress they endure, with women being more affected than men. This phenomenon is more prevalent among younger individuals and those living in disadvantaged areas (6). Conversely, findings from the research suggest that chronic joint pain can impact the quality of sleep in individuals (7). Sleep quality refers to the ability to remain in a state of sleep while exhibiting all the characteristics of а particular stage of sleep (8). Sleep disorders are correlated to specific abnormalities in cognitive function and the processing of emotional information, serving as a primitive sign of mental disorder. These problems may arise from a physical ailment or high levels of stress and depression. It is common in society to see individuals with demanding routines experiencing difficulties with sleep. Persistent sleep disruptions that interfere with daily activities can result in a decline in sleep quality (9).

Given the negative impact of psychosomatic disorders on individuals, it is crucial to focus on their treatment. The results of various studies, such as Galbiati et al. (10), examining the impact of metacognitive therapy on insomnia, and Vandekerckhove and Wang (2018),investigating the significance of emotion regulation on sleep quality, have all confirmed the importance of emotion regulation in enhancing psychological well-being. Strengthening emotion regulation skills can improve individuals' psychological characteristics as it involves being aware of, accepting, and effectively controlling emotions in oneself and others (11).

Emotion regulation, in simpler terms, refers to the skill of managing one's emotions (12). Gross's model (2002) suggests that emotion regulation consists of conscious and unconscious techniques aimed at improving, maintaining, or lessening the emotional, behavioral, and cognitive components of an emotional response (13). The gross model consists of five stages, each with compatible and incompatible strategies. Incompatible strategies provide temporary relief but can cause more harm in the long run (14). The presence of emotion regulation skills helps individuals control impulsive behaviors, such as self-injury, risky actions, and physical aggression, in times of emotional distress (15). Hence, Emotion Regulation Training could be a beneficial method in the holistic treatment of psychosomatic disorders and enhancing the overall quality of life for those affected. To the best of our knowledge, there has been limited research on this topic. Yang et al. (2023) conducted a study on Emotion Regulation Training and its impact on sleep quality (16), while Kraiss et al. (2020) separately confirmed the connection between emotion regulation and psychological well-being (17).

2. Objectives

On the whole, this educational method, which emphasizes emotional regulation and positive coping strategies, assists women in better managing the stress and emotions brought on by their chronic diseases. Improving their ability to identify, understand, and manage their emotions will probably lead to an improvement in their mental well-being. Furthermore, by lessening the stress and anxiety triggered by chronic pain and enhancing coping skills, the quality of sleep may also see an improvement in these patients. Improved sleep quality not only affects the patients' physical well-being but can also help reduce their mental stress and enhance their overall well-being. This study aims to investigate whether utilizing the Gross model for emotion regulation is beneficial for the psychological well-being and sleep quality of women suffering from psychosomatic disorders, such as rheumatic and joint pains, taking into account the current research gap.

3. Methods

The research methodology utilized in this study involved a semi-experimental approach, focusing on pre-test and post-test comparisons with a control group. The study included all women with psychosomatic

disorders in Farahan City who sought treatment at the city hospital for rheumatic or joint pains during the first half of 2022 and were diagnosed with conditions such as rheumatism, arthritis, or pseudo-rheumatism. A total of 40 women with psychosomatic disorders related to rheumatic or joint pains were conveniently selected to participate in the study. The sample size was determined according to the research design, with two groups consisting of 20 individuals each, resulting in a total of 40 participants (Figure 1). The group size included 20 women per group, determined using the G-Power software with an effect size of 0.95, an alpha value of 0.05, and а test power of 0.90.





Individuals eligible to participate in the study were those who were suffering from psychosomatic disorders and currently receiving treatment from a psychiatrist. Additionally, they had to be women aged 18 and above with basic literacy skills in reading and writing. Participants must not have any other mental illnesses in their medical records and must provide consent to take part in the research. The exclusion criteria for the study included missing more than three therapy sessions, a history of substance abuse, lack of motivation to continue participation, presence disorders other of psychiatric than psychosomatic disorders, and mistakes in questionnaires. completing initial Noncompliance with the prescribed exercises and homework after therapy sessions was also grounds for exclusion.

Patients with psychosomatic disorders from Farahan Hospital, Arak City, were chosen based on recommendations from the hospital psychiatrist. After meeting the criteria for participation, 22 out of the initial 67 male patients were excluded through administrative procedures, resulting in a final group of 45 patients. From this group, 3 individuals were taken out of the control group. In order to ensure equality between the two groups, the researcher removed two more participants to

make them identical in number. Following the selection process, the subjects were divided into experimental and control groups using a random and arbitrary method. It is important to note that odd numbers were assigned to the control group, while even numbers were assigned to the experimental group. Participants in both groups completed a pretest before being randomly divided into experimental and control groups. The experimental underwent emotion group regulation intervention sessions based on the Gross model for eight 90-minute sessions. The control group did not receive any intervention. After the experimental period, both groups completed a post-test. The intervention was conducted by the researcher at Farahan City Hospital. An Emotion Regulation Training program consisting of 8 sessions was conducted following the standard protocol developed by Gross (13). Details of the intervention sessions are provided in Table 1.

Table 1. Standard protocol of Gross's Emotion Regulation Training (13)					
Session	Subject	Objectives			
1	Getting to know and communicate with each other	 Establishing relationships among group members and initiating a mutual connection between the group leader (psychologist) and the members. Presenting the primary and secondary objectives of the group and engaging members in discussions about collective and secondary goals. Clarifying the rationale and steps involved in the intervention process. Outlining the structure and guidelines for participation in the group. 			
2	Select position	Providing emotional training			
3	Select position	 Evaluating an individual's emotional vulnerability and emotional skills through following steps: 1.Self-evaluate to recognize one's emotional experiences. 2.Self-evaluate to determine one's level of emotional vulnerability. 3.Self-evaluate to identify strategies for self-regulation. 			
4	Correct position	Making a change in the exciting situation			
5	Expand attention	Teaching skills to change focus			
6	Cognitive assessment	Changing cognitive assessments			
7	Response adjustment	Changing the behavioral and physiological consequences of emotion			
8	Evaluation and application	Re-evaluation and planning for implementing training programs			

Ryff Scale of Psychological Well-Being:

The questionnaire was developed by Ryff in 1989 and updated in 2002 (18). This revised version includes six factors, with the total score of these factors serving as the overall psychological well-being score. The replies to this scale are rated on a 5-point Likert scale, with a higher score indicating better psychological well-being. Items 1, 3, 4, 5, 9, 10, 13, and 17 receive reverse scoring. The correlation between the short version of the Ryff Psychological Well-Being Scale and the original scale has varied between 0.70 and 0.89, as reported (18). Khanjani et al. (2013) confirmed the validity and reliability of the questionnaire in Iran. Their research reported reliability coefficients for the questionnaire's subscales ranging from 0.71 to 0.76 using Cronbach's alpha method, with a total reliability of 0.71 (19). The coefficients for the six factors, namely self-acceptance, environmental mastery, positive relationship with others, purpose in life, personal growth, and autonomy, were found to be 0.51, 0.76, 0.75, 0.52, 0.73, 0.72, respectively, and 0.71 for the entire scale.

Pittsburgh Sleep Quality Index:

This 17-item instrument was designed in 2004 to assess sleep quality and disturbances over the previous month (20). Four of these items are open-ended and not used in statistical analysis. Responses the to questionnaire fall in the range from 1 (not experienced at all) to 4 (three or more times a week), with total scores ranging between 13 (healthy sleep quality) and 52 (impaired sleep quality). Research by Castro et al. (2004) found the questionnaire to have a reliability of 0.83 based on Cronbach's alpha method (20). In Iran, Ghassemzadeh et al. analyzed the psychometric characteristics of this survey. The face validity and reliability of the questionnaire were assessed using the splithalf method, resulting in a Cronbach's alpha coefficient of 0.79 (21).

To conduct MANCOVA and ANCOVA, it was crucial to validate the assumptions of these tests. This included checking the normality of data distribution using the Kolmogorov-Smirnov and Shapiro-Wilk tests. It was also important to ensure that there was no significant difference in the two-group test with one-way ANOVA, assess the sphericity with Bartlett's test, evaluate the linearity with scatter diagrams, and assume the homogeneity of the covariance matrix with the Box-M test. Finally, the homogeneity of variances was confirmed through Levene's test. The researchers used Wilks's lambda multivariate statistic analyze the to

importance of the mean difference in variables related to psychological well-being. Wilks's lambda is a type of multivariate test, along with Pillai's Trace, Roy's Largest Root, and Hotelling's Trace, that helps determine if there is a statistically significant difference between groups in the linear combination of specific subscales.

4. Results

In this study, the control group consisted of 5% of individuals aged 18 to 20, 15% between 20 and 30, 55% in the 30-40 age group, and 25% between 40 and 50. In contrast, the experimental group was formed by 25% of individuals aged between 20 and 30, 30% in the 30-40 age group, 40% between 40 and 50, and 5% over 50 years old. Additionally, considering educational background, 15% of the participants in the control group had a bachelor's degree, 30% a diploma, 25%. On the other hand, the experimental group had 35% below the diploma level, 30% with a diploma, 25% with a bachelor's degree, and 10% with a postgraduate degree.

The data in Table 2 indicates that the experimental group showed an improvement in psychological well-being and its components in the post-test compared to the pre-test. Furthermore, the sleep quality scores of the experimental group decreased in the post-test compared to the pre-test.

The findings of the test are presented in Table 3 (Wilks's lambda=0.209; F_{27.6}=16.988; P<0.05), demonstrating a variance in psychological well-being aspects between the control and experimental groups. According to the eta coefficient, the Gross model's Emotion Regulation Training has a significant effect of 0.79 on enhancing the psychological suffering of women well-being from psychosomatic disorders (F=0.209, P<0.01, η²=0.79, η²=16.988).

The analysis results from Table 4 demonstrate that the Emotion Regulation

Training using the Gross model significantly impacts personal growth in women with psychosomatic disorders, with an effect

size of 0.40. The F value for the mean difference in self-acceptance is 60.73, indicating a significant impact of 0.65 on selfacceptance in this population (F=60.73; P<0.01, n²=0.65, n²=60.73). Furthermore, the adjusted mean scores for personal growth (8.91 and 10.59) and self-acceptance (7.58 and 10.36) in the experimental group are higher compared to the control group, demonstrating the effectiveness of Emotion Regulation Training based on the Gross model in improving psychological well-being among women with psychosomatic disorders.

The information provided in Table 5 demonstrates that there is a significant difference in mean sleep quality due to Emotion Regulation Training using the Gross model (F_{39.1}=17.34; P=0.001). The impact of this training on sleep quality in women with psychosomatic disorders is significant at 0.32 η²=0.32, η²=17.34 (F, (P<0.01. 39.1)). Additionally, the control group had higher mean sleep quality scores (25.73 and 30.22) compared to the experimental group, suggesting that Emotion Regulation Training based on the Gross model effectively enhances sleep quality in women with psychosomatic disorders.

Variable		Cor	itrol	Test	
		Mean	SD	Mean	SD
In daman daman	Pre-test	8.20	2.04	8.15	1.60
Independence	Post-test	Control Te Mean SD Mean Pre-test 8.20 2.04 8.15 Post-test 8.30 2.03 8.55 Pre-test 8.10 1.80 8.55 Post-test 8.70 1.75 10.80 Pre-test 9.25 1.33 9.00 Post-test 8.90 1.37 9.25 Pre-test 10.00 1.92 9.80 Post-test 9.90 1.65 9.90 Pre-test 9.70 1.63 9.55 Post-test 9.45 1.28 9.70 Pre-test 9.45 1.50 10.50 Pre-test 7.45 1.50 10.50 Pre-test 52.20 5.07 52.50 Post-test 52.75 5.95 58.70 Pre-test 37.00 4.24 31.40 Post-test 32.0 4.18 23.90	1.19		
Demonstration of the	Pre-test	8.10	1.80	8.55	2.35
Personal growth	Post-test	8.70	1.75	10.80	1.67
For the second second	Pre-test	9.25	1.33	9.00	1.38
Environmental mastery	Post-test	8.90	1.37	9.25	0.91
	Pre-test	10.00	1.92	9.80	1.24
Positive communication with others	Post-test	9.90	1.65	9.90	1.21
Durmanafiliana	Pre-test	9.70	1.63	9.55	1.64
Purposetuiness	Post-test	Mean SD Mean test 8.20 2.04 8.15 test 8.30 2.03 8.55 test 8.10 1.80 8.55 test 8.70 1.75 10.80 test 9.25 1.33 9.00 1 test 9.00 1.65 9.90 1 test 9.70 1.63 9.55 1 test 9.45 1.28 9.70 1 test 6.95 1.85 7.45 1 test 52.20 5.07 52.50 1 test 52.75 5.95 58.70 1	1.22		
Calf an anti-	Pre-test	6.95	1.85	7.45	1.05
Self-acceptance	Post-test	Mean SD Mean e-test 8.20 2.04 8.15 st-test 8.30 2.03 8.55 e-test 8.10 1.80 8.55 e-test 8.70 1.75 10.80 e-test 9.25 1.33 9.00 e-test 8.90 1.37 9.25 e-test 10.00 1.92 9.80 e-test 9.90 1.65 9.90 e-test 9.70 1.63 9.55 e-test 9.70 1.63 9.55 st-test 9.70 1.63 9.70 e-test 5.2.20 5.07 52.50 st-test 52.75 5.95 58.70 e-test 37.00 4.24 31.40	1.61		
Developing to all the inter-	Pre-test	52.20	5.07	52.50	5.55
Psychological well-being	Post-test	52.75	5.95	58.70	4.81
	Pre-test	37.00	4.24	31.40	2.98
Sleep Quality	Post-test	32.0	2.03 8.55 1.80 8.55 1.75 10.80 1.33 9.00 1.37 9.25 1.92 9.80 1.65 9.90 1.63 9.55 1.28 9.70 1.85 7.45 1.50 10.50 5.95 58.70 4.24 31.40 4.18 23.90	23.90	2.85

Table 2. Descrip	tive statistics	of the scores o	f research varia	bles bv	pre-test and	oost-test
			r cocaron vana		pre cescana	

Table 3. Results of multivariate tests investigating the effect of Emotion Regulation Training

Test	Value	F	d.f	Df	Р	η²
Pillai	0.791**	16.988	6	27	0.001	0.791
Wilks's Lambda	0.209**	16.988	6	27	0.001	0.791
Hotelling	3.775**	16.988	6	27	0.001	0.791
Roy's Largest Root	3.775**	16.988	6	27	0.001	0.791

P<0.001**, P<0.05*

	Variable	Variable SS MS F Sig		Sig	η	
	Independence	0.840	0.840	1.47	0.234	0.044
	Personal growth	26.565	26.565	21.47	0.000	0.402
C	Environmental mastery	SS MS F Sig 0.840 0.840 1.47 0.234 1 26.565 26.565 21.47 0.000 1 1.752 1.752 3.50 0.070 1 with others 0.122 0.122 0.20 0.660 1 with others 0.122 0.122 0.20 0.660 1 73.167 73.167 60.73 0.000 1 18.302 0.572 - - 1 39.589 1.237 - - 1 with others 19.789 0.618 - - 1 with others 19.789 0.618 - - 1 - - 1 - <td>0.099</td>	0.099			
Group	Positive communication with others	0.122	0.122	0.20	0.660	0.006
	Purposefulness	0.718	0.718	1.01	0.322	0.031
	Self-acceptance	73.167	73.167	F Sig 1.47 0.234 21.47 0.000 3.50 0.070 0.20 0.660 1.01 0.322 60.73 0.000	0.655	
	Independence	18.302	0.572			
Purposerumess0.7180.7Self-acceptance73.16773.1Independence18.3020.5Personal growth39.5891.2Environmental mastery16.0110.5Positive communication with others19.7890.6Purposefulness22.7030.7Solf acceptance28.5521.3	1.237					
F	Environmental mastery	16.011	0.500			
Error	Positive communication with others	19.789	0.618			
	Purposefulness	22.703	0.709			
	Self-acceptance	SS IVIS F Sig 0.840 0.840 1.47 0.234 26.565 26.565 21.47 0.000 1.752 1.752 3.50 0.070 h others 0.122 0.122 0.20 0.660 0.718 0.718 1.01 0.322 73.167 73.167 60.73 0.000 18.302 0.572 39.589 1.237 16.011 0.500 h others 19.789 0.618 22.703 0.709 38.553 1.205 105.775 52.775 h others 79.600 59.775 184.975				
	Independence	105.775				
	Personal growth	155.500				
Modified	Environmental mastery	52.775				
Total	Positive communication with others	79.600				
	Purposefulness	59.775				
	Self-acceptance	184.975				

Table 4. ANOVA analysis results to investigate the difference in psychological well-being scores

Table 5. ANOVA analysis results to investigate the difference in sleep quality scores

Variable		SS	df	MS	F	Sig	η
Sleep Quality	Group	124.939	1	124.939	17.34	0.001	0.319
	Error	266.617	37	7.206	-	-	-
	Modified Total	1148.975	39	-	-	-	-

5. Discussion

This research aimed to examine the impact of Emotion Regulation Training using the Gross model on the psychological wellquality being and sleep of women experiencing psychosomatic disorders, such as rheumatic and joint pains. The findings from the analysis indicated that the Gross model-based Emotion Regulation Training had a positive effect on enhancing the psychological well-being of women with psychosomatic disorders. The results of a study conducted by Hasanzadeh Namin et al. (22), Kazemi Rezaei et al. (23), and Kraiss et al. (17) are in agreement with this finding. According to Karbalai et al.'s study (2021), emotion regulation, cognitive flexibility, and mindfulness are positively and significantly related to psychological well-being. However, the suppression component of emotion regulation has a negative and significant relationship with psychological well-being (24). The findings of research by Hasanzadeh Nemin et al. (2018) suggest that positive and negative cognitive strategies for regulating emotions directly and indirectly impact the psychological well-being of female students (22). According to Kazemi Rezaei et al. (2018), training in emotional regulation skills based on dialectical behavior therapy influences the cognitive regulation of emotions and the quality of life among cardiovascular patients (23). Based on the results of a study by Kraiss et al. (2020), there is a connection between emotion regulation and the psychological well-being of individuals with mental disorders (17). It can be inferred from the above explanation that individuals' capacity to regulate their emotions may aid in recognizing their and others' emotions in various scenarios, understanding how emotions influence behavior, and ultimately responding appropriately (25).

Emotion Regulation Training has the potential to benefit individuals with

disorders, where psychosomatic physical symptoms stem from underlying mental health issues. By learning how to regulate their emotions, particularly negative ones, patients may achieve a greater sense of mental serenity and subsequently alleviate their physical and emotional distress (26). Addressing the discomfort and challenges associated with these conditions could ultimately enhance the psychological well-being of women affected by psycho-physical disorders. Furthermore, many individuals grappling with both physical and mental health issues often experience heightened levels of stress and anxiety due to their conditions, underscoring the importance of interventions aimed at restoring emotional promoting equilibrium and psychological healing. Improving patients' capacity to manage pain and difficulties, along with acquiring skills for regulating emotions, can assist them in dealing more effectively with the illness, treatment, and psychological strain that may arise (27). If emotions are not effectively managed, they can impact relationships and overall well-being, leading to potential disruptions in both physical and mental health. Therefore, providing emotional regulation training for individuals with psychosomatic disorders can enhance their control over emotional experiences and how they are expressed, ultimately improving mood, emotions, and psychological well-being through the Gross model (28).

Additionally, the findings from a univariate covariance analysis indicated that Gross model-based Emotion Regulation Training had a 0.32 impact on enhancing the sleep quality of women experiencing psychosomatic disorders. The results of studies by Vandekerckhove et al. (11) support this finding. Similarly, Vanderkercho et al. (2018) discovered that effective emotion regulation could improve sleep quality by reducing negative emotions stemming from daily stress (11). In summary, it can be inferred that poor emotion regulation can result in disorders, strained relationships, and a lack of life skills. Emotional dysregulation patterns that hinder people's competence can lead to suffering from various mental disorders. Conversely, effective and appropriate emotional regulation can contribute significantly to the overall well-being and physical health of individuals (29).

Gross's (2002)model of emotion suggests that strategies regulation for managing emotions play a critical role in how individuals evaluate and respond to emotional situations. Emotion Regulation Training helps individuals learn how to navigate their emotional experiences, express their feelings, choose appropriate coping strategies, and interact with others effectively. This training can help individuals cultivate a sense of calmness challenging mental during Furthermore, emotional situations. dysregulation can result in pain and an inability to cope with emotions (13). Training effective emotional regulation on can decrease anxiety and stress levels. Feelings of anxiety and worry can disrupt sleep patterns. Learning to manage emotions can also help in controlling negative thoughts and images that might interfere with sleep. These negative thoughts are often the root cause of sleep disorders (11). Additionally, the mindfulness techniques taught in emotional regulation training can help create a peaceful state of mind and body, promoting better sleep. As a result, the mental and emotional resilience gained from such training can boost overall mental health and well-being, leading to improved sleep quality (30). Therefore, enhancing emotional control through various psychological and mental strategies could indirectly improve the sleep experiencing quality of women psychosomatic disorders. It can be argued that mishandling negative emotions, such as sorrow, rage, and worry, can negatively impact both physical and mental well-being (11). Recognizing stressors and their impact on individuals can facilitate appropriate responses, ultimately reducing stress levels. Furthermore, emotions play a significant role in the development and perpetuation of mental disorders. Educating individuals on emotional regulation techniques can significantly alleviate symptoms associated with mental disorders (28).

The research had limitations, as it only focused on a sample of 40 women in Farahan psychosomatic disorders Citv with complaining of rheumatic and joint pains in 2022. Therefore, caution should be taken when applying the findings to men and other samples with psychosomatic diseases. It is recommended that similar studies be carried out in various cities and also on male subjects to allow for a broader and more accurate generalization. Furthermore, it is advised to include a follow-up period of 6 or 12 months for research conducted on this topic. Healthcare professionals can also incorporate psychological and emotiontherapies, focused such as emotion regulation based on the Gross model, to enhance the sleep quality of psychosomatic patients alongside medical treatments.

6. Conclusion

managing Research and training in emotions can assist patients in enhancing their emotional well-being and mental health by effectively utilizing their emotions and displaying caring behaviors, leading to improved physical and mental health. Consequently, individuals will likely experience improved sleep quality, better overall physical health, and more stable emotions.

Acknowledgments: The authors express appreciation to all individuals who contributed to the realization of this study. This research was approved by the Research Ethics Committees at Alzahra University (IR.ALZAHRA.REC.1401.191). **Availability of data and materials:** Upon reasonable request to the corresponding author.

Conflicts of interest: The researchers state that the study was carried out without any affiliations or financial interests that could be seen as a possible conflict of interest.

Consent for publication: Not applicable

Ethics approval and consent to participate: The ethics committee at Alzahra Universityhas approved this study (IR.ALZAHRA.REC.1401.191) and it aligns with the Declaration of Helsinki. Subjects have provided their informed consent. Written informed consent was secured from all participants before their involvement in the research.

Financial disclosure: No financial support was received for this study.

Author contributions: M. D. F: Contributed to the conception of the work, revising the draft, approving the final version of the manuscript, and agreeing on all aspects of the work; A. T: Contributed to the data analysis and interpretation, critical revision, and approval of the final version of the manuscript; M. D. F.: Contributed to acquiring data, drafting the manuscript, and approving the final version; A. T. and E. M. S.: Contributed to the conception and design of the study, critical revision, and approval of the manuscript's final version; M. D. F: Contributed to the conception and design of the study, data interpretation, drafting of the manuscript and critical revision, and approval of final version.

References

- Tatayeva R, Ossadchaya E, Sarculova S, Sembayeva Z, Koigeldinova S. Psychosomatic Aspects of The Development of Comorbid Pathology: A Review. Med J Islam Repub Iran. 2022 Dec 12;36:152. doi: 10.47176/mjiri.36.152. PMID: 36636258; PMCID: PMC9826781. <u>https://doi.org/10.47176/mjiri.36.152</u> PMid:36636258 PMCid:PMC9826781
- Torrubia-Pérez E, Reverté-Villarroya S, Fernández-Sáez J, Martorell-Poveda MA. Analysis of Psychosomatic Disorders According to Age and Sex

in a Rural Area: A Population-Based Study. J Pers Med. 2022 Oct 18;12(10):1730. doi: 10.3390/jpm12101730. PMID: 36294869; PMCID: PMC9605088.

https://doi.org/10.3390/jpm12101730 PMid:36294869 PMCid:PMC9605088

- Bahrami Rad M, Rafezi Z. Predicting pain acceptance based on perceived stress and coping strategies in individuals with rheumatoid arthritis. Health Psychology. 2019 Feb 20;7(28):151-66. https://doi.org/10.30473/hpj.2019.38617.3887
- Åslund C, Starrin B, Nilsson KW. Psychosomatic symptoms and low psychological well-being in relation to employment status: the influence of social capital in a large cross-sectional study in Sweden. Int J Equity Health. 2014 Mar 4;13:22. doi: 10.1186/1475-9276-13-22. PMID: 24593256; PMCID: PMC3973887. <u>https://doi.org/10.1186/1475-9276-13-22</u> PMid:24593256 PMCid:PMC3973887
- Muqodas I, Kartadinata S, Nurihsan J, Dahlan T, Yusuf S, Imaddudin A. Psychological well-being: A preliminary study of guidance and counseling services development of preservice teachers in Indonesia. InInternational Conference on Educational Psychology and Pedagogy-" Diversity in Education" (ICEPP 2019) 2020 Feb 7 (pp. 56-60). Atlantis Press

https://doi.org/10.2991/assehr.k.200130.080

- Wyns A, Hendrix J, Lahousse A, De Bruyne E, Nijs J, Godderis L, Polli A. The Biology of Stress Intolerance in Patients with Chronic Pain-State of the Art and Future Directions. J Clin Med. 2023 Mar 14;12(6):2245. doi: 10.3390/jcm12062245. PMID: 36983246; PMCID: PMC10057496. <u>https://doi.org/10.3390/jcm12062245</u> PMid:36983246 PMCid:PMC10057496
- Zambelli Z, Halstead EJ, Fidalgo AR, Dimitriou D. Good Sleep Quality Improves the Relationship Between Pain and Depression Among Individuals With Chronic Pain. Front Psychol. 2021 May 7;12:668930. doi: 10.3389/fpsyg.2021.668930. PMID: 34025533; PMCID: PMC8138032. <u>https://doi.org/10.3389/fpsyg.2021.668930</u> PMid:34025533 PMCid:PMC8138032
- Nelson KL, Davis JE, Corbett CF. Sleep quality: An evolutionary concept analysis. Nurs Forum. 2022 Jan;57(1):144-151. doi: 10.1111/nuf.12659. Epub 2021 Oct 5. PMID: 34610163. https://doi.org/10.1111/nuf.12659 PMid:34610163
- Merrill RM. Mental Health Conditions According to Stress and Sleep Disorders. Int J Environ Res Public Health. 2022 Jun 29;19(13):7957. doi: 10.3390/ijerph19137957. PMID: 35805615; PMCID: PMC9265846.

https://doi.org/10.3390/ijerph19137957 PMid:35805615 PMCid:PMC9265846

- 10. Galbiati A, Sforza M, Scarpellino A, Salibba A, Leitner C, D'Este G, Mombelli S, Ferini-Strambi L, Castronovo V. "Thinking About Thinking" in Insomnia Disorder: The Effect of Cognitive-Behavioral Therapy for Insomnia on Sleep-Related Metacognition. Front Psychol. 2021 9;12:705112. Sep doi: 10.3389/fpsyg.2021.705112. 34566784; PMID: PMCID: PMC8458623. https://doi.org/10.3389/fpsyg.2021.705112 PMid:34566784 PMCid:PMC8458623
- 11. Vandekerckhove M, Wang YL. Emotion, emotion regulation and sleep: An intimate relationship. AIMS Neurosci. 2017 Dec 1;5(1):1-17. doi: 10.3934/Neuroscience.2018.1.1. PMID: 32341948; PMCID: PMC7181893. <u>https://doi.org/10.3934/Neuroscience.2018.1.1</u> PMid:32341948 PMCid:PMC7181893
- 12. Kozubal M, Szuster A, Wielgopolan A. Emotional regulation strategies in daily life: the intensity of emotions and regulation choice. Front Psychol. 2023 Aug 14;14:1218694. doi: 10.3389/fpsyg.2023.1218694. PMID: 37645071; PMCID: PMC10460911. https://doi.org/10.3389/fpsyg.2023.1218694
 PMid:37645071 PMCid:PMC10460911
- 13. Gross JJ. Emotion regulation: affective, cognitive, and social consequences. Psychophysiology. 2002 May;39(3):281-91. doi: 10.1017/s0048577201393198. PMID: 12212647. https://doi.org/10.1017/S0048577201393198 PMid:12212647
- Duggan JM, Toste JR, Heath NL. An examination of the relationship between body image factors and non-suicidal self-injury in young adults: the mediating influence of emotion dysregulation. Psychiatry Res. 2013 Apr 30;206(2-3):256-64. doi: 10.1016/j.psychres.2012.11.016. Epub 2012 Dec 7. PMID: 23219105. https://doi.org/10.1016/j.psychres.2012.11.016

PMid:23219105 15. Hall K, Youssef G, Simpson A, Sloan E, Graeme L,

- Perry N, Moulding R, Baker AL, Beck AK, Staiger PK. An Emotion Regulation and Impulse Control (ERIC) Intervention for Vulnerable Young People: A Multi-Sectoral Pilot Study. Front Psychol. 2021 Apr 1;12:554100. doi: 10.3389/fpsyg.2021.554100. PMID: 33868064; PMCID: PMC8047628. <u>https://doi.org/10.3389/fpsyg.2021.554100</u>PMid:33 868064 PMCid:PMC8047628
- 16. Yang CC, Llamas-Díaz D, Bahena YA, Cabello R, Dahl RE, Magis-Weinberg L. Emotion regulation difficulties and sleep quality in adolescence during the early stages of the COVID-19 lockdown. J Affect Disord. 2023 Oct 1;338:92-99. doi: 10.1016/j.jad.2023.05.036. Epub 2023 May 23. PMID: 37230269; PMCID: PMC10204279.

https://doi.org/10.1016/j.jad.2023.05.036

PMid:37230269 PMCid:PMC10204279

- 17. Kraiss JT, Ten Klooster PM, Moskowitz JT, Bohlmeijer ET. The relationship between emotion regulation and well-being in patients with mental disorders: A meta-analysis. Compr Psychiatry. 2020 Oct;102:152189. doi: 10.1016/j.comppsych.2020.152189. Epub 2020 Jun 18. PMID: 32629064. https://doi.org/10.1016/j.comppsych.2020.152189 PMid:32629064
- 18. Ryff CD. Psychological well-being in adult life. Current directions in psychological science. 1995 Aug;4(4):99-104. <u>https://doi.org/10.1111/1467-8721.ep10772395</u>
- 19. Khanjani M, Shahidi S, Fathabadi J, Mazaheri MA, Shukri A. Factorial structure and psychometric properties of the short form (18 questions) of RIFF psychological well-being scale in male and female students. Thought and behavior in clinical psychology. 2013;32(9):36-27.
- 20. Rey de Castro J, Gallo J, Loureiro H. Cansancio y somnolencia en conductores de ómnibus y accidentes de carretera en el Perú: estudio cuantitativo [Tiredness and sleepiness in bus drivers and road accidents in Peru: a quantitative study]. Rev Panam Salud Publica. 2004 Jul;16(1):11-8. Portuguese. doi: 10.1590/s1020-49892004000700002. PMID: 15333261. https://doi.org/10.1590/S1020-49892004000700002 PMid:15333261
- 20. Ghassemzadeh H, Mojtabai R, Karamghadiri N, Ebrahimkhani N. Psychometric properties of a Persian-language version of the Beck Depression Inventory--Second edition: BDI-II-PERSIAN. Depress Anxiety. 2005;21(4):185-92. doi: 10.1002/da.20070. PMID: 16075452. https://doi.org/10.1002/da.20070 PMid:16075452
- 22. Hasanzadeh Namin F, Peymani J, Ranjbaripoor T, Abolmaali Alhoseini K. (2019). Predictive model of psychological well-being based on resilience considering the mediating role of perceived stress . Journal of Psychological Science. 18(77), 569-578. URL: http://psychologicalscience.ir/article-1-446en.html
- 23. Kazemi Rezaei S V, Kakabraee K, Hosseini S S. The Effectiveness of Emotion Regulation Skill Training Based on Dialectical Behavioral Therapy on Cognitive Emotion Regulation and Quality of Life of Patients With Cardiovascular Diseases. J Arak Uni Med Sci 2019; 22 (4) :98-111<u>https://doi.org/10.32598/JAMS.22.4.90</u> URL: http://jams.arakmu.ac.ir/article-1-6059-en.html

24. Karbalaie M, Yazdanbakhsh K, Karimi P. Predicting Psychological Well-Being Based on Emotion Regulation, Cognitive Flexibility and Mindfulness in Cancer Patients. Health Psychology. 2021 May 22;10(37):141-

60.https://doi.org/10.30473/hpj.2021.50258.4653

- 25. Lane RD, Smith R. Levels of Emotional Awareness: Theory and Measurement of a Socio-Emotional Skill. J Intell. 2021 Aug 19;9(3):42. doi: 10.3390/jintelligence9030042. PMID: 34449662; PMCID: PMC8395748. <u>https://doi.org/10.3390/jintelligence9030042</u> PMid:34449662 PMCid:PMC8395748
- 26. Renna ME, Quintero JM, Fresco DM, Mennin DS. Emotion Regulation Therapy: A Mechanism-Targeted Treatment for Disorders of Distress. Front Psychol. 2017 Feb 6;8:98. doi: 10.3389/fpsyg.2017.00098. PMID: 28220089; PMCID: PMC5292405. <u>https://doi.org/10.3389/fpsyg.2017.00098</u> PMid:28220089 PMCid:PMC5292405
- 27. Hammoudi Halat D, Soltani A, Dalli R, Alsarraj L, Malki A. Understanding and Fostering Mental Health and Well-Being among University Faculty: A Narrative Review. J Clin Med. 2023 Jun 30;12(13):4425. doi: 10.3390/jcm12134425. PMID: 37445459; PMCID: PMC10342374. <u>https://doi.org/10.3390/jcm12134425</u> PMid:37445459 PMCid:PMC10342374
- 28. Menefee DS, Ledoux T, Johnston CA. The Importance of Emotional Regulation in Mental Health. Am J Lifestyle Med. 2022 Jan 12;16(1):28-31. doi: 10.1177/15598276211049771. PMID: 35185423; PMCID: PMC8848120. https://doi.org/10.1177/15598276211049771
 PMid:35185423 PMCid:PMC8848120
- 29. Jauniaux J, Tessier MH, Regueiro S, Chouchou F, Fortin-Côté A, Jackson PL. Emotion regulation of others' positive and negative emotions is related to distinct patterns of heart rate variability and situational empathy. PLoS One. 2020 Dec 31;15(12):e0244427. doi: 10.1371/journal.pone.0244427. PMID: 33382784; PMCID: PMC7774949. https://doi.org/10.1371/journal.pone.0244427
- Sawyer H. Mindfulness: Strategies to implement targeted self-care. J Interprof Educ Pract. 2023 Jun;31:100614. doi: 10.1016/j.xjep.2023.100614. Epub 2023 Feb 13. PMID: 36816353; PMCID: PMC9924360.

https://doi.org/10.1016/j.xjep.2023.100614 PMid:36816353 PMCid:PMC9924360